

Antiox Version



Cutting-Edge Supplement with potent Antioxidant, Anti-inflammatory, Health Body Composition and Performance Effects



Antiox version IV is the most complete and most effective antioxidant formula available on the market today. It combines more than 60 potent natural antioxidants that effectively neutralize free radicals and improve health, body composition, strength, and physical and mental performance.

Antiox version IV - <https://metabolicdiet.com/product/antiox/>. Updated May 3, 2019

The information below on the new Antiox version IV is in near final draft form and will be expanded and revised over the next few months. For now, this latest information will give you the flavor of just what Antiox will do for you in helping you achieve your health, body composition and performance goals.

Antioxidants form a front-line defense against cell damage caused by free radicals, which are involved in muscle, joint and tendon damage and inflammation, degenerative arthritis and even in the aging process. The use of antioxidants can reduce free radical damage that occurs when we exercise and can also attenuate the ongoing damage to injured tissues caused by free radicals, thus accelerating the healing process.

Antioxidants, such as vitamins C and E (see under Vitamins below), selenium, green tea, reduced glutathione and N-acetyl-cysteine (NAC) can play an important role in reducing inflammation and fatigue, decreasing tissue damage, and in both preventing and treating injuries.

Various antioxidants, such as vitamin E, have been found to be useful in the treatment of some forms of arthritis¹ and in dealing with the oxidative stress of exercise.² As well, oxidative damage has been shown to contribute to the pathogenesis of injuries and arthritis, and the use of antioxidants, such as NAC,³ shown to have therapeutic value for reducing endothelial dysfunction, inflammation, fibrosis, invasion and cartilage erosion.

A study found that a combination of 2 antioxidants, selenomethionine and epigallocatechin-gallate (the main antioxidant in green tea extract), had beneficial effects on catabolic and anabolic gene

expression of articular chondrocytes.⁴ The authors of the study concluded that “Our data provide insights into the mechanisms whereby ECGg and selenium modulate chondrocyte metabolism. Despite their differential mechanisms of action, the 2 compounds may exert global beneficial effects on articular cartilage.”

Antiox version IV Nutrition Panel

Supplement Facts:		Serving Size: 6 Tablets			
		Servings Per Container: 30			
	Amount Per Serving	% Daily Value			
	Amount Per Serving	% Daily Value			
Vitamin A (Beta Carotene)	10,000 IU	200%	Alpha Lipoic Acid	100 mg	*
Vitamin C (Ascorbic Acid)	400 mg	667%	Turmeric Root Extract	100 mg	*
Vitamin E (d-Alpha Tocopherol Acetate)	200 IU	667%	Panax Ginseng Root Extract	100 mg	*
Vitamin B1 (Thiamin HCL)	10 mg	666%	Cranberry Juice Powder	100 mg	*
Vitamin B2 (Riboflavin)	10 mg	588%	Rosemary Leaf Extract	100 mg	*
Niacinamide	25 mg	125%	Superoxide Dismutase	62.5 mg	*
Vitamin B6 (Pyridoxine HCL)	10 mg	500%	Ferulic Acid	60 mg	*
Folic Acid	400 mcg	100%	Cinnamon Bark Extract	60 mg	*
Vitamin B12 (Cyanocobalamin)	100 mcg	1667%	Schizandra Berry Extract	60 mg	*
Biotin	1000 mcg	333%	L-Glutamine	60 mg	*
d-Calcium Pantothenate	10 mg	100%	Pomegranate Fruit Extract	60 mg	*
Calcium (Calcium Phosphate)	250 mg	25%	Betaine HCL	50 mg	*
Magnesium (Citrate)	200 mg	50%	Grape Seed Extract	50 mg	*
Zinc (Optizinc – Zinc monomethionine)	15 mg	100%	L-Carnosine	50 mg	*
Selenium (Sodium Selenite and Selenomethionine)	50 mcg	70%	Rutin	50 mg	*
Copper (Gluconate)	200 mcg	10%	Bilberry Fruit Extract	40 mg	*
Manganese (Sulfate)	5 mg	250%	Hawthorn Berry Extract	30 mg	*
Bioperme (Piper nigrum) (fruit)	5 mg	*	Ginger Root Powder	30 mg	*
L-Taurine	250 mg	*	Pine Bark Extract (<i>Pinus strobus</i>)	25 mg	*
L-Methionine	200 mg	*	Phosphatidylserine Complex	20 mg	*
Ginkgo Biloba Leaf Extract	200 mg	*	Lutein	5 mg	*
Grape Skin Extract	200 mg	*	Lycopene	5 mg	*
Green Tea Extract (<i>Camellia sinensis</i>) (leaf)	200 mg	*	Antiox™ Proprietary Complex 816.5 mg		
Creatine Monohydrate	200 mg	*	Citrus Bioflavonoids Complex, Garlic Bulb Powder (<i>Allium sativum</i>),		
Eleuthero Root Extract	200 mg	*	Omega 3 Fish Oil Powder, L-Carnitine, Coenzyme Q10, Glutathione (Reduced)		
Quercetin Dihydrate	120 mg	*	L-Cysteine HCL, Resveratrol, Astaxanthin, Zeaxanthin		
Milk Thistle Seed Extract	100 mg	*			
N-Acetyl Cysteine	100 mg	*			
Other Ingredients: Cellulose, Stearic Acid, Magnesium Stearate, Hypromellose, Modified Cellulose Gum, Silicon Dioxide					
*Daily Value Not Established					

Antiox version IV

Antiox is so much more than just the usual antioxidant supplement. Because of the mix of ingredients, it also has other beneficial effects including decreasing inflammation, protecting muscle from microtears, apoptosis of muscle fibers, and damage from pro-oxidants stemming from oxidative phosphorylation and ATP formation, improving recovery, and helping to improve body composition and exercise/sports performance.

All you have to do is look at the nutritional supplement panel to see how truly unique Antiox is and how much it can do to prevent the counter-productive aspects of exercise while enhancing exercise's beneficial effects.

Antiox contains the several antioxidants, including vitamin C, beta-carotene, vitamin E, zinc, pycnogenol, resveratrol, astaxanthin, zeaxanthin, various amino acids and various herbs. However, it also contains many other antioxidants that give it a full rounded effect without an overemphasis on just a few isolated antioxidants and contains other ingredients that facilitate the formation of endogenous antioxidants as well as complement the antioxidant and health benefits of Antiox.

For example, various nutrients are necessary for priming the endogenous antioxidant systems including some minerals as copper, zinc and selenium (all present in Antiox) that contribute to the antioxidant defense system by acting as co-factors for antioxidant Cu-Zn superoxide dismutase and glutathione peroxidase activities.

Overall, Antiox contains over 60 nutrients that have antioxidant activity and/or increase endogenous antioxidant activity. As well, many of these ingredients have other properties that result in beneficial effects on health, energy metabolism, body composition and performance.

Having this large mix of antioxidants, as well as plant and herbal extracts, and supporting nutrients, makes Antiox not only more natural than other antioxidants on the market but also makes it more effective and bypasses the problems associated with using just a few antioxidants by themselves.

Studies have shown that a complex mixture of ingredients, including plant and herbal extracts, are synergistic in their effects on metabolism and their health and performance benefits. For example, a review paper published in 2017 concluded that herbal synergism often found in polyherbal formulations were found effective to combat disease heterogeneity as found in complex pathogenesis of Alzheimer's disease.⁵

As such, Antiox, with its individual antioxidants as well as full and partial plant extracts, and fruit complexes, provides the most complete full spectrum antioxidant effects one would expect from a combination of single antioxidants and natural extracts from herbs, plants, and fruits. In fact, I've made Antiox more like a super antioxidant and anti-inflammatory food rather than a nutritional supplement since it contains so many ingredients, not just a mix of the most effective antioxidants but also many complex herbal, and plant extracts.

Antioxidants Enhance the Anabolic Effects of Exercise

Before I continue with information on oxidative stress and antioxidants, I want to make one thing clear - antioxidants as supplied by Antiox does not impair the anabolic effects of exercise.

Exercise damages skeletal muscle, with the extent of the damage depending on the intensity and duration of resistance, aerobic and sports specific exercise. The damage to muscle is caused by both mechanical and oxidative stress.

While some people feel that the use of antioxidant supplements are counter-productive in that they decrease the exercise induced mechanical muscle injury and more importantly the subsequent inflammatory response, that leads to a response by the body to repair the damage and increase muscle hypertrophy, this is not the case.

Many studies have shown that the inflammatory response secondary to high mechanical muscle fiber stimulation after intense exercise mediates the initial steps involved in regeneration, healing, and muscle hyperplasia and hypertrophy.

However, studies have shown that antioxidants do not effect muscle fiber stimulation secondary to exercise, which leads to a beneficial effect on muscle mass and strength, **but can decrease excessive muscle fiber damage and oxidative stress, which are counter-productive and not needed for the hypertrophic and performance enhancing response to exercise.**^{6,7,8} As such, it is unlikely that the use of antioxidants by athletes will impair the repair, hypertrophic, and performance enhancing response secondary to counter-productive muscle fiber damage and microtears in the musculoskeletal system that can lead to injuries and decrease the beneficial effects of exercise.

On the other hand, the use of antioxidants does affect some counter-productive processes that occur in the body secondary to intense and/or prolonged, exhaustive exercise, including fibrosis, mitochondrial damage, muscle fiber cell apoptosis, irreparable structural damage, and immune system dysfunction, which inhibit the ability of muscle to hypertrophy, and decrease both short and long term exercise performance.

Antiox has ingredients that reduce inflammation and severe muscle damage although productive muscle damage secondary to exercise is not unduly affected so that the secondary anabolic response is not diminished but in fact due to other ingredients is increased. In other words, Antiox decreases oxidative damage and inflammation without blocking the beneficial cellular adaptation to exercise.

More on this below under the Ergogenic Effects of Antioxidants.

In the next section, I'll cover some of the ingredients in Antiox and what they're all about. After that I'll go into details about oxidative stress, antioxidants and other information that you'll find useful in understanding oxidative stress and antioxidants.

Some of the Ingredients in Antiox version IV

There are a variety of antioxidants that are commercially available, including vitamins C and E, selenium, beta-carotene, glutathione, superoxide dismutase, gamma oryzanol, coenzyme Q10, cysteine, and histidine dipeptides (such as carnosine). Antiox contains over 60 ingredients with antioxidant and anti-inflammatory activity. We'll discuss some of these below.

Glutathione

Glutathione (gamma-glutamylcysteinylglycine) is one of the most effective endogenous and exogenous antioxidants. In both in rats and humans it has been found that exhaustive physical exercise causes a change in glutathione redox status in blood and that antioxidant administration, i.e., oral vitamin C, N-acetyl-L-cysteine, or glutathione, is effective in preventing oxidation of the blood glutathione pool after physical exercise in rats⁹.

In one study, the changes of glutathione status, oxidative injury, and antioxidant enzyme systems after an exhaustive bout of treadmill running and/or hydroperoxide injection in male rats were studied¹⁰. In this study, concentrations of total and reduced glutathione in muscle were significantly increased after exhaustive exercise. Concentrations of glutathione-related amino acids glutamate, cysteine, and aspartate were also significantly increased. As well, endogenous antioxidant enzymes such as glutathione peroxidase, glutathione reductase, superoxide dismutase, and catalase activities were significantly elevated in muscle after exhaustive exercise. Overall, this study shows that exhaustive exercise can impose a severe oxidative stress on skeletal muscle and that glutathione systems and antioxidant enzymes are important in coping with free radical-mediated muscle injury.

Glutathione, a potent endogenous antioxidant that protects major organs from oxidant injury, may not be present in adequate amounts during intense exercise. Since glutathione reserves may be increased with supplements, nutrition strategies that include the provision of GSH monoester may lend additional support to musculoskeletal and other tissues that are at risk for injury from oxygen free radicals during catabolic states¹¹.

N-Acetylcysteine

N-acetyl-L-cysteine (or NAC) is the N-acetyl derivative of L-cysteine (a non-essential amino acid) and a reduced thiol donor. As with other amino acids, only L-isomers are biologically active. While Antiox also contains smaller amounts of cysteine, NAC is thus the preferred form for oral use for systemic effects while cysteine is more useful for use by the gastrointestinal tract.¹²

Cysteine serves as a component of coenzyme A which is involved in carbohydrate and fat metabolism. Cysteine is also a precursor of taurine. Numerous studies have shown the clinical use of NAC supplements in chronic¹³ but apparently not acute¹⁴ pulmonary problems. NAC is marketed in European and other countries for use in chronic pulmonary diseases.

NAC has been shown to have hepatoprotectant effects against various liver toxic compounds such as alcohol, acetaminophen, carbon tetrachloride, and drugs used in cancer chemotherapy.¹⁵ Much of the protective effects are due to NAC's antioxidant properties and its ability to increase cellular levels of glutathione.

As well, NAC and other reducing agents have been shown to be useful for the treatment of AIDS and the wasting seen in AIDS patients since they have been shown to decrease viral replication and to inhibit the action of inflammatory cytokines.^{16,17}

Aside from protein synthesis, cysteine/NAC is an essential component of glutathione (L-gamma-glutamyl-L-cysteinylglycine - one of the main cellular antioxidants). NAC and cysteine also act as potent antioxidants both by themselves¹⁸ or in concert with glutathione and other antioxidants.¹⁹

NAC, because of its potent antioxidant properties, may prove useful in decreasing the oxidative stress seen with exercise.²⁰ Human studies performed by Dr. Lester Packer of the University of California at Berkeley, and presented at the 1996 American College of Nutrition meeting in San Francisco found that supplementation with NAC improved exercise performance times and reduced oxidative stress by preserving glutathione levels during exercise.

Since oxidative stress is considered a causal factor in human muscle fatigue and overtraining,²¹ this finding is extremely important because it provides evidence of positive effects of amino acid supplementation on human exercise performance and recovery. If this response were to continue for an adequate time period, then enhanced results from resistance training (faster increase in muscle mass and strength) could be expected.

Studies in the last decade have shown that the most promising thiol donors are L-cysteine, NAC, glutathione and alpha lipoic acid, have significant effects on exercise performance.^{22, 23, 24, 25, 26, 27, 28, 29}

Of these the prototypical compound in this category is N-acetylcysteine (NAC), a reduced thiol donor with antioxidant properties, which has effects both directly and indirectly by inhibiting glutathione oxidation and supporting reduced glutathione synthesis and resynthesis.

NAC infusion in well trained individuals increased each of muscle NAC, total and reduced glutathione, cysteine and cystine, and substantially enhanced performance during prolonged exercise.³⁰ Oral use of NAC has recently been demonstrated to enhanced handgrip performance.³¹

A recent study looked at the possible interaction between ROS and Na⁺,K⁺-pump activity in skeletal muscle, and a possible role in fatigue during exercise.³² The results showed that the antioxidant NAC attenuates muscle fatigue, in part via improved K⁺ regulation and attenuating the decline in Na⁺,K⁺ pump activity, and point to a role for ROS in muscle fatigue.

In fact the effects of NAC on fatigue was examined in a paper in which the author felt that taking NAC may constitute doping. ****

Selenium

Selenium is an essential trace element that provides antioxidant protection in concert with vitamin E (both in Antiox). Several selenoproteins have been identified, but only one, glutathione peroxidase (SeGSHpx), a selenoenzyme, has a known function, that of neutralizing toxic lipid hydroperoxides³³. Selenium, however has several metabolic effects not associated with glutathione peroxidase³⁴.

Many of the nutrients have multiple actions in the body. For example, selenium also has significant anti-inflammatory properties and involved in thyroid health and function.³⁵³⁶

Vitamin C

Vitamin C is essential to proper collagen synthesis, and this is evident in the vitamin C deficiency disease scurvy, in which the collagen fibers synthesized in the body cannot form fibers properly, resulting in lesions, blood vessel fragility and poor wound healing.

Vitamin C has been shown to have some anticatabolic effects that likely involves decreasing exercise induced cortisol but may also have some effects through its antioxidant action. Conversely, some of the anticatabolic effects of antioxidants may be mediated through a decrease in cortisol.

Antioxidants may be of some use in training induced muscle ischemia and injury. Research shows that exercise can adversely affect muscle tissue by increasing the formation of free radicals. These free radicals can then lead to muscle fatigue, inflammation and muscular damage.³⁷ During normal conditions free radicals are generated at a low rate and neutralized by antioxidant enzymes in the liver and skeletal muscle and other systems. Unfortunately, the increase in free radicals caused by exercise accompanies a simultaneous decrease in the supply of antioxidants to handle them. Vitamin E, for instance, can be severely decreased by training thus depleting muscle of its major antioxidant force.³⁸

A study examined the potential protective effect of pretreatment with corticosteroids or antioxidants (ascorbic acid or allopurinol) in rabbits with reperfusion-induced damage to skeletal muscle after ischemia.³⁹ In this study 4 hours of limb ischemia induced by a pneumatic tourniquet, followed by reperfusion for 1 hour, caused a considerable amount of ultrastructural damage to the anterior tibialis muscles accompanied by a rise in circulating creatine kinase activity. Pretreatment of animals with depomedrone by a single 8 mg bolus injection led to a preservation of the anterior tibialis structure on both light and electron microscopy. High-dose continuous intravenous infusion with ascorbic acid (80 mg/hr) throughout the period of ischemia and reperfusion also preserved skeletal muscle structure, although allopurinol in various doses had no protective effect.

These data are fully compatible with a mechanism of ischemia/reperfusion-induced injury to skeletal muscle, involving generation of oxygen radicals and neutrophil sequestration and activation. They also indicate that damage to human skeletal muscle caused by prolonged use of a tourniquet is likely to be reduced by simple pharmacological interventions.

Vitamin C is necessary for collagen synthesis and is a strong antioxidant with beneficial effects on pro-inflammatory cytokines.⁴⁰ Research on vitamin C shows that it may have important effects in reducing pain and inflammation secondary to exercise. In one study 400 mg daily of vitamin C reduced post exercise pain and inflammation.⁴¹ Vitamin C is involved in the enzymatic hydroxylation of proline to form 4-hydroxyproline, an amino acid that is an integral part of collagen and elastin.

Vitamin E

One of the most effective and central antioxidants is alpha-tocopherol (vitamin E). It appears to have an important role as both a primary antioxidant and as a compound that enhances the effect of both endogenous and exogenous antioxidants.

Vitamin E, a lipophilic chain-breaking antioxidant which inhibits lipid peroxidation in isolated mitochondrial membranes and protects membranes from oxidative damage, performs an antioxidant role in biological membranes by acting as a one-electron reductant⁴². The primary oxidation product of vitamin E is the tocopheroxyl radical. Reduction of the tocopheroxyl radical can occur by reactions with water-soluble anti-oxidants such as ascorbate or glutathione, resulting in the recycling of vitamin E⁴³.

It has been shown that free radical-mediated lipid peroxidation and glutathione depletion may be involved in the toxic manifestations of various compounds⁴⁴. It has also been shown that vitamin E may largely modulate the expression of toxicity by glutathione depleting agents. Experiments carried

out with vitamin E deficient or supplemented diets indicate that the pathological phenomena occurring as a consequence of glutathione depletion depend on hepatic levels of vitamin E⁴⁵.

According to one study, deficiency in vitamin E may be related to arterial lesions⁴⁶. Vitamin E can mitigate some of the adverse effects of fat in diets. The lipid peroxidation caused in the liver, heart and intestine by a diet containing autoxidized oil was suppressed when the same tocopherol level as fresh soybean oil was added to the diet⁴⁷. Uptake of oxidatively modified low-density lipoprotein (LDL) by cells in the arterial wall is believed to be an important early event in the development of atherosclerosis.

This lipid peroxidation mechanism, which can readily be rationalized by the known chemistry of the alpha -tocopheroxyl radical and by the radical isolating properties of fine emulsions such as LDL, explains how reagents which reduce the alpha -tocopheroxyl radical (i.e. vitamin C and ubiquinol-10) strongly inhibit lipid peroxidation in vitamin E-containing LDL⁴⁸.

Level of plasma tocopherol (vitamin E) which has free radical scavenging properties rises significantly during intensive exercise⁴⁹. Mobilization of tocopherol could help to prevent lipoperoxidation phenomena occurring in exercising skeletal muscle. A combination of vitamin E and coenzyme Q10 (both in Antiox) was found to attenuate exercise induced lipid peroxidation and muscular damage.⁵⁰

Green Tea Extract

The constituents of green tea are polyphenolic compounds termed catechins. The most abundant catechin in green tea is (–)-epigallocatechin 3-gallate (EGCG) although others are also present in lesser quantities.

Green tea extract, besides being rich in antioxidants, also has significant effects on inflammation and the musculoskeletal system.^{51, 52, 53} For example green tea catechins have been shown to inhibit inflammation and cartilage degradation, and have therapeutic effects on a variety of musculoskeletal problems including arthritis.^{54, 55, 56, 57, 58,}

Green tea extract may well prove to be more useful than green tea itself. A recent study found that green tea extract supplements retain the beneficial effects of green and black tea and allow larger doses of tea polyphenols to be used without the side effects of caffeine associated with green and black tea beverages.⁵⁹

Rutin and Quercetin

These two flavonoids have been shown in studies to have significant anti-inflammatory activity in cases of both acute and chronic inflammation.⁶⁰ A review article concluded that there is evidence to suggest that flavonoids may be beneficial to connective tissue for several reasons, which include the limiting of inflammation and associated tissue degradation, the improvement of local circulation, as well as the promoting of a strong collagen matrix.⁶¹ These compounds also have significant antioxidant properties.

Quercetin may have properties that downregulate or inhibit cyclooxygenase-2 safely.⁶² As well, studies have shown that quercetin has beneficial effects on exercise, especially on exercise induced muscle damage.^{63,64} A recent study found that **“Fourteen days of Q supplementation seems able**

to attenuate the severity of muscle weakness caused by eccentric-induced myofibrillar disruption and sarcolemmal action potential propagation impairment.”⁶⁵

Turmeric

Curcumin, the active component of turmeric, is documented to have anti-inflammatory and antioxidative benefits.⁶⁶ As an antioxidant, curcumin reduces the activity of certain enzymes, inhibiting all branches of the arachidonic acid cascade. Thereby, these plant extracts reduce inflammation.

Turmeric exhibits marked anti-inflammatory action and has been shown to be as effective as some anti-inflammatory drugs. For example, in a double-blinded trial, post-surgical patients receiving curcumin experienced reductions in stiffness and joint swelling comparable to the effects of phenylbutazone, a potent anti-inflammatory drug.⁶⁷

Of all the spices and herbal preparations it seems that only the spice turmeric has any anti-inflammatory effects. This was the conclusion of a study of a variety of Ayurvedic and herbal preparations, which was presented recently at the 9th Asia Pacific League of Associations for Rheumatology Congress.

In this study, a variety of herbal and Ayurvedic preparations were tested in rats. The rats were fed oral doses of the varied herbal and Ayurvedic recipes. Only turmeric showed anti-inflammatory effects when tested on irritated paws of the rats.

Several studies have shown the effectiveness of curcumin, especially when coupled with piperine which increases absorption of curcumin (both are in Antiox) on exercise induced muscle damage and soreness, and recovery as well as on improving body composition and exercise performance.
68697071727374 In one study the combination of curcumin and piperine resulted in an improvement of in sprint mean power output 24 hours post-exercise.⁷⁵

As well, other studies have shown the value of curcumin in the prevention and treatment of neurological dysfunction such as Alzheimer's disease and other neurological diseases.⁷⁶⁷⁷

Resveratrol

Resveratrol is a potent anti-oxidant with significant anti-aromatase activity.⁷⁸⁷⁹ Researchers have shown that resveratrol may have significant anti-aging effect and extend lifespan and healthspan.⁸⁰

As well, resveratrol has been shown to decrease fatigue. enhances body composition and both physical and mental performance, and improves testicular function.⁸¹⁸²⁸³⁸⁴⁸⁵⁸⁶⁸⁷⁸⁸⁸⁹⁹⁰⁹¹

Studies have also shown that resveratrol, and the B vitamins in Antiox, can counteract some of the detrimental effects of environmental pollution especially first, second, and third hand cigarette smoke and air pollution.⁹²⁹³⁹⁴

Both the ingredient resveratrol and grape seed extract that contains resveratrol and many other beneficial ingredients, are in Antiox.

Carnosine

Carnosine, a dipeptide made up of the amino acids alanine and histidine (histidyl-alanine or beta-alanyl-L-histidine) was added to Antiox because of its many beneficial effects. It has been shown to have significant antioxidant, anti-inflammatory, anti-ischemic properties, increases healing, enhances the immune system, improves skeletal muscle function, and provides health and anti-aging effects.^{95,96,97,98,99,100,101,102,103,104,105,,106,107,108,109,110}

It also inhibits glycation, a destructive protein/sugar reaction that occurs in the body and which contributes to aging through a number of mechanisms including the breakdown of connective tissue, a loss of elasticity, and a decrease in cellular hydration. Carnosine, along with alpha lipoic acid, provides protection against glycation and premature aging.

Carnosine also works well with other ingredients in Antiox. For example carnosine, chromium and cinnamon and polyphenols, all in Antiox, have shown beneficial effects on body composition and insulin sensitivity.¹¹¹¹¹²

Coenzyme Q10 (ubiquinone-10)

Coenzyme Q10 acts as an electron carrier of the respiratory chain in mitochondria. Under the various forms of stress and inflammation, demand for coenzyme Q10 increases which must be met by dietary intake in order to optimize mitochondrial function.

As well, it has been shown that the reduced form of coenzyme Q10 is an important physiological lipid-soluble antioxidant that scavenges free radicals generated chemically within liposomal membranes.^{113,114} It has also been shown that it reduces oxidative stress associated with strenuous exercise in rats, healthy adults and young athletes.¹¹⁵¹¹⁶¹¹⁷¹¹⁸ As noted above, vitamin E and ubiquinone increase physical working capacity of experimental animals.¹¹⁹

Generation of free radicals and subsequent lipid peroxidation have been proposed to contribute to delayed tissue damage. One study has found that ascorbate and ubiquinol levels were decreased after trauma.¹²⁰ In this study, changes in tissue levels of ubiquinol, but not ascorbate reflected the degree of trauma. The authors suggest that ubiquinol levels may provide a useful marker of the oxidative component of the secondary injury response.

Beta carotene

Beta carotene is the most popularly known carotenoid, although over 600 carotenoids from natural sources that have been characterized. Of these less than 60 can be converted to vitamin A. Many dietary carotenoids, both with and without provitamin A activity, are found in the blood and tissues of humans. Carotenoids, however, have biological actions apart from their function as precursors of vitamin A.

Studies have shown that particular carotenoids enhance the immune system, inhibit mutagenesis, reduce nuclear damage, protect against photo-induced tissue damage, and quench highly reactive singlet oxygen under certain conditions and can block free radical-mediated reactions¹²¹.

Although Beta-Carotene is only one of the nutritionally active carotenoid (making up 15-30% of total serum carotenoids) it is often the only carotenoid considered in antioxidant regimens. Using only

synthetic beta carotene without any of the associated carotenoids, although theoretically beneficial may show some adverse effects. For example, in epidemiological studies, the intake of carotenoids from natural sources such as certain fruits and vegetables has been associated with a decreased incidence of certain cancers especially lung cancer.

The main problem with epidemiological associations is that because they do not show cause and effect, it is difficult to properly interpret the data and draw proper conclusions. For example while one study shows that dietary beta carotene and vitamin E supplements reduce the risk of lung cancer in nonsmoking men and women¹²², another study shows that the use of beta carotene in smokers may actually increase the incidence of lung cancer.

The synthetic all-trans isomer of beta carotene was recently shown to possess antioxidant properties towards the formation of oxidized low density lipoprotein. The all-trans isomer of beta -carotene is more effective in inhibiting the susceptibility of lipoproteins to lipid peroxidation and in reducing the cellular uptake of oxidized LDL by macrophages¹²³.

Taurine

Taurine, a sulfur-containing amino acid and the second most abundant amino acid, and the most abundant free amino acid, found in skeletal muscle tissue, has many properties that can enhance the training effect, including its abilities to increase growth hormone, protect joints, and protect the liver, as well as its antioxidant and anabolic effects. Taurine has also been shown to have insulin like effects and to help control cell volume. The volumizing effect on muscle cells is felt to lead to an increase in protein synthesis.

Over the years oral taurine administration has been shown to help muscle cramping in patients with liver cirrhosis and myotonic dystrophy. Several studies have suggested that it may also help to alleviate muscle cramps occurring during and after exercise. But there's more to the story.

One study on rats has shown that oral taurine supplementation may increase muscle performance and reduce muscle injury caused by exercise.¹²⁴ The aim of the study was to determine if increasing muscle levels of taurine would decrease free radical damage after exercise-induced injury. The authors found that first of all taurine levels rose in muscle after supplementation, and secondly that running performance was improved by the taurine supplementation. Thus it appears taurine supplementation may facilitate exercise performance and reduce some of the counterproductive muscle injury caused by exercise.

Another study also in rats showed that taurine is useful for reducing physical fatigue and muscle damage during exercise training, presumably due to its antioxidant effects and the beneficial effects that taurine has on metabolism and on muscle and cardiac functions.¹²⁵

And yet another study found that taurine may attenuate exercise-induced DNA damage and enhance the capacity of exercise due to its cellular protective properties.¹²⁶ Taurine, along with carnosine has also shown protective effects on testicular injury.¹²⁷

And as if that weren't enough, there is some evidence to show that taurine may enhance training further by decreasing training induced fatigue. One paper has shown that Na⁺-K⁺-ATPase activity is depressed with fatigue, regardless of training state, suggesting that this may be an important determinant of fatigue.¹²⁸ Another paper associated fatigue and training with reduced Ca²⁺-ATPase activity.¹²⁹ Previous studies have shown that taurine stimulates Na⁺-K⁺-ATPase activity and also the pumping rate of the Ca²⁺-activated ATPase pump.

While the evidence is circumstantial at present and a direct study linking the two research findings needs to be done, it's quite possible that supplemental taurine, besides all the positive effects we know it has, may also reduce exercise induced fatigue.

On a preventative level, taurine administration has been shown to increase taurine concentrations in skeletal muscles, reduce the decrease in taurine concentration in skeletal muscles on exercise, and up-regulates physical endurance.¹³⁰

For more information on taurine have a look at my updated 2019 article at https://metabolicdiet.com/wp-content/uploads/2017/product_pdf/Taurine.pdf.

Alpha Lipoic Acid

Alpha lipoic acid (ALA) has potent antioxidant properties intrinsically and secondary to its ability to increase levels of intra-cellular glutathione, and its ability to recycle other antioxidants such as vitamin C, vitamin E and glutathione.^{131,132,133,134,135} ALA and glutathione have been shown to have significant effects in decreasing mercury toxicity in the body.¹³⁶

Alpha lipoic acid also has significant anti-inflammatory properties and has been shown to inhibit IL-1, a proinflammatory cytokine and also inhibit the synthesis of PGE2 by inhibiting COX-2 activity.¹³⁷

This latter mode of action, along with similar effects from CLA, simulates the anti-inflammatory effects of the present class of NSAIDS such as Celebrex, Advil, Aleve, etc. As well, Antiox contains fish oil containing DHA and EPA, which has also been shown to have effects similar to the anti-inflammatory prescription and OTC drugs.¹³⁸

ALA has other beneficial effects, such as decreasing both the pro-inflammatory cytokines, decreasing secondary cortisol elevations, and improving the lipid profile and cardiovascular risk factors.
139140141142

It has been shown to inhibit cross-linking among proteins, a process that contributes to the aging process in the body and especially in collagen-heavy tissues such as skin. Alpha-lipoic acid activates a collagen-regulating factor known as AP-1 that turns on enzymes that digest glycation-damaged collagen and thus make the skin more supple and youthful looking.

Besides having potent antioxidant and anti-inflammatory effects, ALA also has significant anabolic and body composition effects secondary to its beneficial effects on insulin sensitivity and growth hormone and IGF-I secretion, all factors involved in maintaining, repairing and regenerating musculoskeletal tissues.^{143,144,145,146147}

ALA is also useful in reversing mitochondrial dysfunction, especially in aging mitochondria.^{148, 149}

Zinc

Exercise can lead to an increased need for certain nutrients. For example, one study found that there is an increase in selenium requirements with exercise.¹⁵⁰ Problems can arise from exercise induced mineral loss, which is further enhanced by the finding that many of us don't consume adequate amounts of many essential minerals.

Studies have shown that many athletes, and female athletes, in particular, consume diets that have been found to be inadequate for certain key minerals such as zinc, magnesium, copper, and iron. The combination of strenuous exercise and compromised mineral status ultimately leads to low endurance capacity, depressed immune function, and the development of a variety of disease conditions.

One study looked at the effects of zinc deficiency on physical performance and found that low dietary zinc was associated with impaired cardiorespiratory function and impaired metabolic responses during exercise.¹⁵¹

Zinc deficiency in humans is widespread¹⁵² and athletes may be particularly prone to lower plasma zinc levels.¹⁵³ Zinc is a constituent of more than a hundred fundamentally important enzymes, so zinc deficiency has many negative effects on almost every body function.¹⁵⁴ As well, zinc deficiency can adversely affect the reproductive hormones and as such impair athletic efforts.¹⁵⁵

Zinc deficiency adversely affects protein synthesis. In one study the effects of zinc deficiency in rats, on the levels of free amino acid in urine, plasma and skin extract were investigated.¹⁵⁶ Zinc deficiency adversely affected skin protein synthesis. Especially where a deficiency may be present, supplemental zinc has resulted in an increase the secretion of growth hormone and IGF-I,¹⁵⁷ and testosterone¹⁵⁸ and to raise plasma testosterone and sperm count.^{159,160}

A study looking at the effects of zinc supplementation on wrestlers found that the results obtained at the end of the study indicate that zinc supplementation (as well as several other ingredients in Antiox including NAC and ALA) prevents production of free radicals by **activating the endogenous antioxidant system**.¹⁶¹ This activation is important as it coincides with the effects of exercise, which also activates the endogenous antioxidant system and leads to endogenous antioxidants that enhance the beneficial effects of exercise on body composition and performance. The authors concluded that “physiologic doses of zinc supplementation to athletes may beneficially contribute to their health and performance.”

It's been shown that there is an improvement in insulin resistance with **zinc** supplementation and that zinc is involved in controlling some of the aspects of obesity.¹⁶² Zinc also improves calcium metabolism and thus the beneficial effects that calcium has on fat metabolism.

Magnesium

Magnesium, besides complementing the effects of calcium on obesity¹⁶³ and other functions, also has important effects on its own. Magnesium is involved in numerous processes that affect muscle function including oxygen uptake, energy production and electrolyte balance. Low levels of magnesium promote inflammation^{164,165} and impact on the body's ability to handle stress.¹⁶⁶ These functions are useful in alleviating the release of pro-inflammatory cytokines, and decreasing both insulin resistance and inappropriate cortisol secretion.

There is evidence that marginal magnesium deficiency impairs exercise performance and increases oxidative stress. As well, strenuous exercise increases urinary and sweat losses that may increase magnesium requirements.¹⁶⁷

Recent surveys have shown that a significant number of individuals are magnesium deficient based on their intake. Athletes in sports with weight classes are especially vulnerable to magnesium deficiency due to their weight loss practices. As such, in these athletes, and others who are

magnesium deficient or whose levels are marginal, magnesium supplementation would have beneficial effects on exercise performance.

A recent study found that magnesium supplementation improved alactic anaerobic metabolism, even though the athletes were not magnesium-deficient.¹⁶⁸ Another study found that magnesium supplementation increased strength performance.¹⁶⁹ As well, magnesium may prove effective for muscle cramps.¹⁷⁰

Magnesium has been shown to influence testosterone levels as well as the anabolic peptide IGF-1.¹⁷¹¹⁷² As well, magnesium has been shown to work along with zinc and B6 (both of which are present in Antiox) to produce a significant anabolic effect.¹⁷³

Other Antioxidants

Many other compounds, some touted as ergogenic aids, have shown significant antioxidant properties.

Gamma oryzanol, another compound widely used by athletes, while not having any significant anabolic effects, does have some antioxidant activity¹⁷⁴.

Cysteine (decreases cross linkage of proteins), **taurine**¹⁷⁵ (may have a protective effect in joints by decreasing the degradation of hyaluronic acid), histidine dipeptides¹⁷⁶, pantothenic acid, copper, zinc, selenium.

Cinnamon has been shown to have significant antioxidant, anti-inflammatory, and anti-obesity effects. A recent study found that the use of cinnamon results in fewer molecules involved in the body's fat-storing process and more antioxidant and anti-inflammatory molecules that protect the body from the damages of stress.¹⁷⁷

Antioxidant Interactions

Many antioxidant compounds act in concert with both other exogenous compounds and with endogenous antioxidants. For example ubiquinol-10 spares alpha-tocopherol when both antioxidants are present in the same liposomal membranes and that ubiquinol-10, like alpha-tocopherol, does not interact with reduced glutathione¹⁷⁸.

Generation of free radicals and subsequent lipid peroxidation have been proposed to contribute to delayed tissue damage. One study has found that ascorbate and ubiquinol levels were decreased after trauma¹⁷⁹. In this study, changes in tissue levels of ubiquinol, but not ascorbate reflected the degree of trauma. The authors suggest that ubiquinol levels may provide a useful marker of the oxidative component of the secondary injury response.

In some cases antioxidants act synergistically. For example water based antioxidants such as Vit C can act in concert with lipophilic antioxidants such as Vit E. In one study Vit E and Vit C acted synergistically to inhibit the oxidation of human low density lipoprotein (LDL)¹⁸⁰. In another study, the combination of vitamins E and C produced a protective effect on parinaric acid peroxidation exceeding the sum of their individual contributions¹⁸¹.

Many antioxidants have a vitamin E-sparing action. The water-soluble antioxidant vitamin C can reduce tocopheroxyl radicals directly or indirectly and thus support the antioxidant activity of vitamin E^{182,183}. In one study, vitamin C and other compounds (including ascorbyl palmitate, propyl gallate, butylated hydroxytoluene, hydroquinone and glutathione) blocked the oxidation of platelet tocopherol. In this same study, it was shown that the use of exogenous natural and/or synthetic antioxidants can prevent the oxidation of endogenous antioxidants¹⁸⁴.

Oxidative Stress

Oxidative stress occurs when the generation of reactive oxygen species (ROS) in a system exceeds the system's ability to neutralize and eliminate them. This condition can arise from a lack of antioxidant capacity and/or an excessive production of ROS.

The end result is an excess of ROS that in turn can damage cellular lipids, proteins and DNA, and causing problems with cellular function. As such, oxidative stress may contribute to a number of diseases and in the aging process.¹⁸⁵

Free Radicals and Tissue Damage

In the past few decades there has been an intense amount of investigation on the role of free radicals in disease and aging, and on the role of antioxidants on attenuating the damaging effects of free radicals.

Free radicals are highly reactive molecules with unpaired electrons. These radicals play an important part in many physiological processes, including the metabolism of food and the utilization of energy substrates during exercise. These free radicals arise from endogenous metabolic processes and increase with physical (such as exercise) and emotional stress. Free radicals can also arise in the body as a result of exposure to environmental pollutants¹⁸⁶¹⁸⁷¹⁸⁸.

The most common free radicals are the superoxide anion and the free radicals derived from its interaction with other molecules. Superoxide radicals are generated from cellular respiratory-chain oxygen use and by a number of other enzyme systems.¹⁸⁹¹⁹⁰ It has been proposed that the superoxide radical (along with its derivatives, such as singlet oxygen and the hydroperoxyl, hydroxyl, and alkoxy radicals) and other free radicals initiate the peroxidation of membrane lipids, which results in cellular damage¹⁹¹.

Also it is felt that the peroxidation of low-density lipoproteins (LDL) and their subsequent cytotoxicity is involved in atherogenesis. One study has shown that there may be a synergistic effect of HDL and antioxidant molecules (such as vitamin E and catechin) for the protection of cells against oxidized LDL, and for the preventative modification of LDL¹⁹². As well, there is evidence to show that over time there is substantial oxidant-induced DNA damage¹⁹³.

In reaction to free-radical reactions and lipid peroxidation that occur in living cells, highly efficient enzymatic and nonenzymatic systems for scavenging free radicals have evolved. The antioxidant enzymatic system consists largely of superoxide dismutase (SOD), catalase, and glutathione peroxidase, which detoxify, respectively, the superoxide radical, hydrogen peroxide, and lipid hydroperoxides.¹⁹⁴¹⁹⁵ In addition to these enzymes, various free-radical scavengers are present, such as the micronutrient selenium, which is a component of glutathione peroxidase, ubiquinols (reduced coenzyme Q), ascorbate (vitamin C), and alpha-tocopherol (vitamin E), cysteine, glutathione, and possibly uric acid.¹⁹⁶¹⁹⁷

It has been postulated that free radicals that are produced during normal metabolism react with cellular components, leading to molecular damage and eventually to the death of vital cells and the aging and death of the organism¹⁹⁸. For example, the free radical theory of aging¹⁹⁹ states that endogenous free radical reactions cause or contribute significantly to: cancer, atherosclerosis, hypertension, Alzheimer's disease, amyloidosis, immune deficiency, arthritis, diabetes and Parkinson's Disease, among others. Proponents of this free-radical theory have advocated the use of various antioxidants in the diet to interrupt free-radical chain reactions and thus prevent or attenuate cellular damage and aging.²⁰⁰

Antioxidants

Antioxidants are compounds that prevent or limit the formation of ROS and their effects on biological structures. Antioxidant defense involves multiple strategies, enzymatic or non-enzymatic.

Antioxidants function to prevent or minimize peroxidative damage in biological systems. Studies have suggested that adequate antioxidant defense can protect the body from the high free radical concentrations that are unavoidable at the present time²⁰¹.

Non-enzymatic compounds include tocopherols, carotenes, ascorbate, glutathione, ubiquinol, and flavonoids. As well, there are the micronutrient elements such as selenium and zinc which are important as integral constituents of protective enzymes via special amino acids (e.g. selenomethionine) or structural components (e.g. Zn-metallothionein).²⁰² Overall, these low-molecular-mass antioxidant molecules add to the enzymatic defense system provided by superoxide dismutases, catalase, thioredoxin reductases, and glutathione peroxidases.

The Ergogenic Effects of Antioxidants

Studies have shown conflicting results on the use of antioxidant supplements by athletes and anyone interested in improving body composition and performance. The problem lies not so much in the studies themselves but in the study parameters. The studies using a small number of antioxidants, especially vitamins C and E, and studies looking at exercise done in moderation, will usually result in the conclusion that the use of antioxidants are counter-productive for improving body composition and performance.²⁰³²⁰⁴

They reach this conclusion because the protocols do not take into account the duality of exercise. Practiced in moderation, exercise increases the production of endogenous antioxidants that are capable of handling the damaging effects of exercise and oxidative stress on metabolism and the musculoskeletal system. Under these conditions the use of exogenous antioxidants can be counter-productive or productive depending the genetic and epigenetic makeup of the athlete.

However strenuous exercise and exercise under less than optimal conditions, causes cell damage and oxidative stress that often overwhelms the endogenous antioxidants and results in damaged muscle, loss of muscle cells, fibrotic changes within skeletal muscle, and cumulative microtears and other changes that increase the possibility of injuries and overtraining.

Antioxidants form a front-line defense against cell damage caused by free radicals, which are involved in muscle, joint and tendon damage and inflammation, degenerative arthritis and even in the aging process. The use of antioxidants can reduce free radical damage that occurs when we

exercise²⁰⁵ and can also attenuate the ongoing damage to injured tissues caused by free radicals, thus accelerating the healing process.²⁰⁶ As well, antioxidants have been shown to enhance aerobic performance.²⁰⁷

Antioxidants, such as vitamins C and E (see under Vitamins below), selenium, green tea, reduced glutathione and N-acetyl-cysteine (NAC) can play an important role in reducing inflammation and fatigue, decreasing tissue damage, and in both preventing and treating injuries.

Various antioxidants, such as vitamin E, have been found to be useful in the treatment of some forms of arthritis²⁰⁸ and in dealing with the oxidative stress of exercise.²⁰⁹ As well, oxidative damage has been shown to contribute to the pathogenesis of injuries and arthritis, and the use of antioxidants, such as NAC,²¹⁰ shown to have therapeutic value for reducing endothelial dysfunction, inflammation, fibrosis, invasion and cartilage erosion.

One study found that a combination of 2 antioxidants, selenomethionine and epigallocatechin-gallate (the main antioxidant in green tea extract), had beneficial effects on catabolic and anabolic gene expression of articular chondrocytes.²¹¹ The authors of the study concluded that “Our data provide insights into the mechanisms whereby ECGg and selenium modulate chondrocyte metabolism. Despite their differential mechanisms of action, the 2 compounds may exert global beneficial effects on articular cartilage.”

Supplementing endogenous antioxidants with supplementation is especially important in intense exercise which can overwhelm the intracellular antioxidant systems and can result in damage to the musculoskeletal and other systems in the body. For example, oxidative stress, the result of the overwhelming of the endogenous antioxidant system, can lead to damage to the reproductive system and decrease sex hormone production over the long term.^{212,213}

Supplemental exogenous antioxidants interact with endogenous antioxidant to provide protection from the increase in free radicals produced by exercise.

Exogenous supplementation may be useful in all forms of exercise, or they may actually be counterproductive in that they may impair the natural increase in endogenous antioxidants. If the exercise intensity is only mild to moderate supplemental antioxidants may, by decreasing the stimulus to enhance the endogenous antioxidant system, be somewhat counterproductive.

However, supplemental antioxidants are crucial in those that perform acute and chronic intense and/or exhaustive exercise and training since this intensity of training produces excessive free radical production and irreparable oxidative damage that overwhelms the endogenous antioxidant system resulting in irreparable tissue damage, an increase proneness to injury, and ill health.

So while skeletal muscle has an amazing capacity to adapt and repair itself, significantly increased levels of physical activity (intense training, chronic long duration exercise and overtraining) exposes the limitation of adaptation to tissue damage and will lead to some degree of maladaptation and musculoskeletal tissue changes that are counter-productive to both skeletal muscle function and performance.

As well, various nutrients are necessary for priming the endogenous antioxidant systems. For example, some minerals as copper, zinc and selenium (all present in Antiox) contribute to the antioxidant defense system by acting as co-factors for antioxidant Cu-Zn superoxide dismutase and glutathione peroxidase activities.

But Antiox contains much more than the usual antioxidants. It includes a mix of antioxidants and other supporting nutrients that makes other antioxidant supplements pale in comparison.

For example, Antiox contains quercetin, which has been shown to increase brain and muscle mitochondrial biogenesis and exercise tolerance, and thus may enhance athletic performance.²¹⁴

Antiox also contains alpha lipoic acid (see above), which has been shown to enhance insulin sensitivity, decrease post exercise lactic acid concentrations, and have potent antioxidant and anti-inflammatory properties.²¹⁵

Overall, Antiox contains over 60 nutrients that have antioxidant activity and/or increase endogenous antioxidant activity. As well, many of these ingredients have other properties that result in beneficial effects on health, energy metabolism, body composition and performance.

Free Radicals and Exercise

Oxygen consumption greatly increases during exercise, which leads to increased free radical production. The body counters the increase in free radical production through the antioxidant defense system. When free radical production exceeds clearance oxidative damage occurs. Free radicals formed during chronic exercise may exceed the protective capacity of the antioxidant defense system, thereby making the body more immune to disease and injury.

Antioxidants have been shown to be useful in many different conditions. Certain preparations have been postulated to retard the aging process by reducing the peroxidation of fatty acids secondary to enhancing the antioxidant enzymes to inhibit free radical activity. Although one such study showed significant effects of certain compounds on some physical symptoms, levels of endogenous antioxidants and hormonal levels (especially serum testosterone) in elderly men²¹⁶, little is known about the use of such compounds and other antioxidants in retarding the aging process.

Data from many studies show that free radicals play important roles in carcinogenesis^{217,218} (in a role as natural inhibitors which may act on cancer initiation or promotion) and aging. It has also been shown that antioxidants act as protective agents in carcinogenesis²¹⁹, protect against heart disease^{220,221}, and may attenuate the effects of aging²²². Administration of antioxidants (vitamins C, E and A) has been found useful for the prevention of postoperative complications²²³.

Studies have shown that antioxidative treatment by vitamins could be an important regimen in the reduction of reperfusion damage. In three separate studies, inhibition of lipid peroxidation by antioxidants improved the function of the transplanted kidney²²⁴, liver²²⁵, and heart²²⁶.

Experimental evidence suggests that oxidative stress due to the accumulation of free radicals plays a role in the pathogenesis of cataracts and that the process can be prevented or ameliorated by antioxidants, such as Vit C and E^{227,228,229,230}. Low antioxidant levels may also be a risk factor for rheumatoid arthritis²³¹, and myocardial infarction²³².

A study has shown that oxygen-derived free radicals contribute to myocardial injury after cardioplegic arrest and that antioxidant therapy should improve myocardial protection²³³. Another study found that low plasma vitamins E and C seems to lead to increased risk of angina in Scottish men²³⁴. Low plasma vitamin E predisposes to angina, and smoking may increase the risk of angina by lowering plasma vitamin C levels in Scottish men.

There is a growing amount of data available on the physiological and pathological effects of free radicals on the musculoskeletal system. It has been shown that exercise can adversely affect muscle tissue by increasing the formation of free radicals, leading to lipid peroxidation and tissue damage.

Exercise results in free radical formation and muscle damage and inflammation. Muscular contraction produces reactive oxygen intermediates that contribute to muscular fatigue²³⁵. Skeletal muscle oxygen utilization greatly increases during high intensity exercise along with an increased rate of ATP utilization that exceeds the rate of ATP generation. The subsequent metabolic stress results in a markedly enhanced rate of production of oxygen free radicals from semiquinone and xanthine oxidase.

Studies have shown that training causes a substantial decrease in vitamin E concentration in the proliferating muscle mitochondrial membranes, thus depleting muscle mitochondria of their major lipid antioxidant²³⁶. Since vitamin E is the major cellular, lipid-soluble, chain-breaking antioxidant, these findings indicate increased free radical reactions in the tissues of exercising animals.

A study found that high altitude exposure adversely affects glutathione metabolism and antioxidant defense mechanisms and these changes can be ameliorated through supplementation of NAC and vitamin E.²³⁷ Another study found that NAC prevents impairment to exposure to hypobaric hypoxia.²³⁸

During normal conditions free radicals are generated at a low rate and subsequently taken care of by the well-developed scavenger and antioxidant systems. It has been shown that exercise training imposes an oxidative stress to the body and induces increases in skeletal muscle antioxidant enzymes particularly glutathione peroxidase^{239, 240}. It is thought that free radical damage to proteins accelerates overall protein catabolism. However, in most circumstances it is felt that most cells are in possession of sufficient antioxidants to control protein damage, so that changing radical fluxes do not greatly change intracellular protein turnover rates²⁴¹.

Thus under normal exercise conditions antioxidant enzymes in liver and skeletal muscle are capable of adapting to exercise to minimize oxidative injury caused by free radicals²⁴². However, a greatly increased rate of free radical production may exceed the capacity of the cellular defense system. Consequently, a substantial attack of free radicals on the cell membranes may lead to a loss of cell viability and to cell necrosis and could initiate the skeletal muscle damage and inflammation caused by exhaustive exercise²⁴³.

It is possible that therapies based on anti-oxidant supply may be valuable in these cases. For example in the thyrotoxic myopathy and cardiomyopathy, which are the major complications of hyperthyroidism, hyperthyroid muscular tissues may undergo several biochemical changes that predispose them to free radical-mediated injury²⁴⁴.

It has been shown that vitamin E protects against thyroxine-induced acceleration of lipid peroxidation in cardiac and skeletal muscles in rats²⁴⁵. The results of this study also suggest that vitamin E protects against lipid peroxidation in hyperthyroid heart and skeletal muscle independently of the changes in oxidative enzymes and antioxidant enzymes.

Exercise training seems to increase muscle antioxidant enzymes and reduced glutathione²⁴⁶. Many reports show that vitamin E levels decrease in the muscle and increase in plasma during exercise. Studies of vitamin E deficiency and supplementation strongly suggest that this vitamin is of protective value during exercise.

It's well known that muscle fatigue with exercise is caused at least in part by oxidative stress and that the use of antioxidants can delay the muscle fatigue. Oxidants produced by exercise depress muscular force by decreasing the contractile response leading to premature fatigue.

In one study, it was shown that free radicals appeared to contribute to fatigue in oxidative skeletal muscle²⁴⁷. As well animal studies have shown that the use of compounds that neutralize free radicals can enhance aerobic performance.²⁴⁸

Several antioxidants, and especially **NAC** have been shown to decrease fatigue and have a positive effect on endurance.²⁴⁹ For example, two recent studies found that NAC attenuates fatigue during whole body exercise²⁵⁰ and during hand grip exercise.²⁵¹

A recent study concluded that **“natural supplementation with Pycnogenol, with proper hydration, good training and nutritional attention may improve training and performances both in normal subjects and in semi-professional athletes performing at high levels in difficult, high-stress sports such as the triathlon.”**²⁵²

Peroxidation of myocardial-membrane phospholipid is considered an important pathogenic component of heart muscle damage in ischemia and reperfusion. Membrane alpha-tocopherol (vitamin E) in the heart can modulate such damage and protect against it²⁵³. There is evidence that vitamins C and E and beta-carotene may protect against lipid peroxidation damage in muscle tissue. In one study use of these three compounds led to reduced CPK and LDH levels, indicating a decrease in muscle tissue damage.²⁵⁴

It would seem logical to assume that as a result of the vast amount of information now available about antioxidants that they would lessen tissue damage and speed recovery and thus might in the long term lead to increased performance. Antioxidants may be useful for athletes by reducing the destructive effects of free radicals on skeletal muscle as well as other tissues and organs, especially the liver and the central and peripheral nervous systems. Increasing levels of antioxidants prior to intensive exercise may have significant effects in reducing muscle injury and catabolism.

It has been shown that antioxidants might be useful in the biological preparation of endurance athletes. One study has demonstrated the antioxidant effects of a new compound, containing selenium, vitamin E, glutathione and cysteine, prepared as tablets²⁵⁵. In this study 3 weeks of antioxidant treatment induced significant increase of the nonproteic -SH and significant decrease of the lipid peroxides in comparison with placebo treatment. In another study the use of an antioxidant vitamin mixture (ascorbate, alpha -tocopherol, and beta -carotene) lowered markers of lipid peroxidation at rest and after exercise²⁵⁶.

Several human and animal studies suggest that strenuous exercise may promote free radical production, leading to lipid peroxidation and tissue damage. On the other hand, there is evidence that vitamins C and E and beta-carotene may protect against such damage. Thus, concordance between the health benefits of exercise and nutrition and a compensatory role of antioxidant nutrients against the potentially harmful effects of exercise suggests that nutrition and exercise should form important components of any regimen for prevention of chronic diseases and/or promotion of optimal health²⁵⁷.

Result suggests the involvement of the oxygen free radicals generated from the xanthine oxidase pathway in the pathogenesis of the ischemic injury of the rat brain²⁵⁸. Xanthine oxidase, which catalyzes hypoxanthine to xanthine and xanthine to uric acid as the last steps of ATP degradation. These reactions catalyzed by xanthine oxidase are considered to be a source of free radicals and may play important roles in the pathogenesis of cerebral ischemic injury²⁵⁹.

The existence of a xanthine oxidase system in skeletal muscle could produce favorable conditions for oxy-radical formation through hypoxanthine degradation, which may contribute to the known muscle tissue injury²⁶⁰.

It is possible to decrease the peroxidation of skeletal muscle-membrane phospholipid and thus decrease the oxidant radical portion of muscle injury and disease processes. In one recent study, the spontaneous lipid peroxidation in liver, kidney and heart decreased with increasing levels of dietary antioxidants²⁶¹.

Results of some studies have shown that some degree of autoxidation occurs in trauma. It appears that in blunt trauma the synthesis and regeneration capacity of glutathione are intact but that the levels of the consumable antioxidants, ascorbic acid, and alpha-tocopherol are compromised²⁶². It is conceivable that the use of antioxidants might decrease the repercussions of musculoskeletal injuries or the microtrauma associated with overtraining.

Antioxidants form a front line defense against cell damage caused by free radicals, which are involved in muscle, joint and tendon damage and inflammation, degenerative arthritis and even in the aging process. The use of antioxidants can reduce free radical damage that occurs when we exercise and can also attenuate the ongoing damage to injured tissues caused by free radicals, thus accelerating the healing process.

Antioxidants, such as vitamins C and E, selenium, green tea, reduced glutathione, N-acetyl-cysteine (NAC), and many others can play an important role in reducing inflammation and fatigue, decreasing tissue damage, and in both preventing and treating injuries.

Various antioxidants, such as vitamin E, have been found to be useful in the treatment of some forms of arthritis²⁶³ and in dealing with the oxidative stress of exercise.²⁶⁴ As well, oxidative damage has been shown to contribute to the pathogenesis of injuries and arthritis, and the use of antioxidants, such as NAC,²⁶⁵ shown to have therapeutic value for reducing endothelial dysfunction, inflammation, fibrosis, invasion and cartilage erosion.

A study found that a combination of 2 antioxidants, selenomethionine and epigallocatechin-gallate (the main antioxidant in green tea extract), had beneficial effects on catabolic and anabolic gene expression of articular chondrocytes.²⁶⁶ The authors of the study concluded that "Our data provide insights into the mechanisms whereby ECGg and selenium modulate chondrocyte metabolism. Despite their differential mechanisms of action, the 2 compounds may exert global beneficial effects on articular cartilage."

Why Studies Vary on Their Findings

A scan of the literature shows a variety of studies that seem at odds on the effects of antioxidants on health, body composition, training, and performance. There are several reasons these discrepancies, including the use of only one or two antioxidants at one time (vitamin C and vitamin E are the most common), differences in subjects used (for example those who exercise for fitness or fun versus those who train hard including but not exclusive to competitive bodybuilders and power athletes, and most elite athletes in any sport), and a multitude of differences in protocols.

Also, few studies take into account possible gene polymorphisms and epigenetic influences that can affect the degree of oxidative stress and damage, and the effects of antioxidants on the response to exercise.^{267, 268}

The bottom line is that if you're serious about your training, whether for personal or competitive reasons, Antiox will give you anabolic, body composition, performance, recovery, injury protection, and other benefits that will help you reach your goals.

References:

- ¹ Sangha O, Stucki G. Vitamin E in the treatment of rheumatic diseases. *Zeitschrift für Rheumatologie* 1998;57(4):207-214.
- ² Sackeck JM, Blumberg JB. Role of vitamin E and oxidative stress in exercise. *Nutrition* 2001;(10):809-14.
- ³ Zafarullah M, Li WQ, Sylvester J, Ahmad M. Molecular mechanisms of N-acetylcysteine actions. *Cell Mol Life Sci* 2003;60(1):6-20.
- ⁴ Andriamanalijaona R, Kypriotou M, Bauge C, Renard E, Legendre F, Raoudi M, Boumediene K, Gatto H, Monginoux P, Pujol JP. Comparative effects of 2 antioxidants, selenomethionine and epigallocatechin-gallate, on catabolic and anabolic gene expression of articular chondrocytes. *J Rheumatol.* 2005;32(10):1958-67.
- ⁵⁵ Dey A, Bhattacharya R, Mukherjee A, Pandey DK. Natural products against Alzheimer's disease: Pharmaco-therapeutics and biotechnological interventions. *Biotechnol Adv.* 2017 Mar - Apr;35(2):178-216.
- ⁶ Piercy RJ, Hinchcliff KW, DiSilvestro RA, Reinhart GA, Baskin CR, Hayek MG, Burr JR, Swenson RA. Effect of dietary supplements containing antioxidants on attenuation of muscle damage in exercising sled dogs. 2000 Nov;61(11):1438-45.
- ⁷ Nakhostin-Roohi B, Babaei P, Rahmani-Nia F, Bohlooli S. Effect of vitamin C supplementation on lipid peroxidation, muscle damage and inflammation after 30-min exercise at 75% VO₂max. *J Sports Med Phys Fitness.* 2008 Jun;48(2):217-24.
- ⁸ Vats P, Singh VK, Singh SN, Singh SB. Glutathione metabolism under high-altitude stress and effect of antioxidant supplementation. *Aviat Space Environ Med.* 2008 Dec;79(12):1106-11.
- ⁹ Sastre-J; Asensi-M; Gasco-E; Pallardo-FV; Ferrero-JA; Furukawa-T; Vina-J. Exhaustive physical exercise causes oxidation of glutathione status in blood: Prevention by antioxidant administration. *AM-J-PHYSIOL-REGUL-INTEGR-COMP-PHYSIOL.* 263/5 32-5 (R992-R995) 1992.
- ¹⁰ Ji LL; Fu R. Responses of glutathione system and antioxidant enzymes to exhaustive exercise and hydroperoxide. *J Appl Physiol* 1992, 72 (2) p549-54.
- ¹¹ Parenteral glutathione monoester enhances tissue antioxidant stores. Robinson-MK; Ahn-MS; Rounds-JD; Cook-JA; Jacobs-DO; Wilmore-DW. *J-PARENTER-ENTER-NUTR.* 16/5 (413-418) 1992.
- ¹² Meister A. Intracellular cysteine and glutathione delivery systems. *Journal of the American College of Nutrition* 1986;5:137-151, .
- ¹³ Ziment I. Acetyl cysteine: a drug with an interesting past and a fascinating future. *Respiration* 1986;50(suppl):26-30.
- ¹⁴ Konrad F, Schoenberg MH, Wiedmann H, Kilian J, Georgieff M. [The application of n-acetylcysteine as an antioxidant and mucolytic in mechanical ventilation in intensive care patients. A prospective, randomized, placebo-controlled, double-blind study]. *Anaesthesist* 1995;44(9):651-8.
- ¹⁵ Smilkstein M, Knapp GL, Kulig KW, et al. Efficacy of oral N-acetyl cysteine in the treatment of acetaminophen overdose. *The New England Journal of Medicine* 1988;319:1557-1562.
- ¹⁶ Roederer M, Ela SW, Staal FJ, Herzenberg LA, Herzenberg LA. N-acetylcysteine: a new approach to anti-HIV therapy. *AIDS Res Hum Retroviruses.* 1992;8(2):209-17.
- ¹⁷ Kalebic T, Schein PS. Organic thiophosphate WR-151327 suppresses expression of HIV in chronically infected cells. *AIDS Research & Human Retroviruses* 1994;10(6):727-33.

-
- ¹⁸ Izzotti A, D'Agostini F, Bagnasco M, Scatolini L, Rovida A, Balansky RM, Cesarone CF, De Flora S. Chemoprevention of carcinogen-DNA adducts and chronic degenerative diseases. *Cancer Research* 1994;54(7 Suppl):1994s-1998s.
- ¹⁹ Bongers V, de Jong J, Steen I, et al. Antioxidant-related parameters in patients treated for cancer chemoprevention with N-acetylcysteine. *European Journal of Cancer* 1995;31A(6):921-3.
- ²⁰ Sen CK, Rankinen T, Vaisanen S, Rauramaa R. Oxidative stress after human exercise: effect of N-acetylcysteine supplementation. *Journal of Applied Physiology* 1994;76(6):2570-7.
- ²¹ Finaud J, Lac G, Filaire E. Oxidative stress : relationship with exercise and training. *Sports Med.* 2006;36(4):327-58.
- ²² Reid MB, Stokic DS, Koch SM, Khawli FA, Leis AA. N-acetylcysteine inhibits muscle fatigue in humans. *J Clin Invest.* 1994;94(6):2468-74. Sen CK, Atalay M, Hanninen O. Exercise-induced oxidative stress: glutathione supplementation and deficiency. *J Appl Physiol.* 1994;77(5):2177-87.
- ²³ Kinscherf R, Hack V, Fischbach T, Friedmann B, Weiss C, Edler L, Bartsch P, Droge W. Low plasma glutamine in combination with high glutamate levels indicate risk for loss of body cell mass in healthy individuals: the effect of N-acetyl-cysteine. *J Mol Med.* 1996;74(7):393-400.
- ²⁴ Medved I, Brown MJ, Bjorksten AR, Leppik JA, Sostaric S, McKenna MJ. N-acetylcysteine infusion alters blood redox status but not time to fatigue during intense exercise in humans. *J Appl Physiol.* 2003;94(4):1572-82.
- ²⁵ Sen CK, Packer L. Thiol homeostasis and supplements in physical exercise. *Am J Clin Nutr.* 2000;72(2 Suppl):653S-69S.
- ²⁶ Vassilakopoulos T, Karatza MH, Katsaounou P, Kollintza A, Zakynthinos S, Roussos C. Antioxidants attenuate the plasma cytokine response to exercise in humans. *J Appl Physiol.* 2003;94(3):1025-32.
- ²⁷ Medved I, Brown MJ, Bjorksten AR, McKenna MJ. Effects of intravenous N-acetylcysteine infusion on time to fatigue and potassium regulation during prolonged cycling exercise. *J Appl Physiol.* 2004;96(1):211-7.
- ²⁸ Hauer K, Hildebrandt W, Sehl Y, Edler L, Oster P, Droge W. Improvement in muscular performance and decrease in tumor necrosis factor level in old age after antioxidant treatment. *J Mol Med.* 2003;81(2):118-25.
- ²⁹ Quadriatero J, Hoffman-Goetz L. N-Acetyl-L-cysteine prevents exercise-induced intestinal lymphocyte apoptosis by maintaining intracellular glutathione levels and reducing mitochondrial membrane depolarization. *Biochem Biophys Res Commun.* 2004 2;319(3):894-901.
- ³⁰ Medved I, Brown MJ, Bjorksten AR, Murphy KT, Petersen AC, Sostaric S, Gong X, McKenna MJ. N-acetylcysteine enhances muscle cysteine and glutathione availability and attenuates fatigue during prolonged exercise in endurance-trained individuals. *J Appl Physiol.* 2004;97(4):1477-85.
- ³¹ Matuszczak Y, Farid M, Jones J, Lansdowne S, Smith MA, Taylor AA, Reid MB. Effects of N-acetylcysteine on glutathione oxidation and fatigue during handgrip exercise. *Muscle Nerve.* 2005;32(5):633-8.
- ³² McKenna MJ, Medved I, Goodman CA, Brown MJ, Bjorksten AR, Murphy KT, Petersen AC, Sostaric S, Gong X. N-acetylcysteine attenuates the decline in muscle Na⁺,K⁺-pump activity and delays fatigue during prolonged exercise in humans. *J Physiol.* 2006 1;576(Pt 1):279-88.
- ³³ Selenium: clinical significance and analytical concepts. Lockitch-G. *Crit-Rev-Clin-Lab-Sci.* 1989; 27(6): 483-541.
- ³⁴ Burk RF. Recent developments in trace element metabolism and function: newer roles of selenium in nutrition. *J Nutr;* 1989 Jul; 119(7); P 1051-4.
- ³⁵ Duntas LH. Selenium and inflammation: underlying anti-inflammatory mechanisms. *Horm Metab Res.* 2009 Jun;41(6):443-7.
- ³⁶ Köhrle J. Selenium and the thyroid. *Curr Opin Endocrinol Diabetes Obes.* 2015 Oct;22(5):392-401.
- ³⁷ Sjodin B, Hellsten Westing Y, Apple FS. Biochemical mechanisms for oxygen free radical formation during exercise. *Sports Med* 1990;10(4):236-54.
- ³⁸ Gohil K, Rothfuss L, Lang J, Packer L. Effect of exercise training on tissue vitamin E and ubiquinone content. *J Appl Physiol* 1987;63(4):1638-41.

-
- ³⁹ Bushell A, Klenerman L, Davies H, Grierson I, Jackson MJ. Ischemia-reperfusion-induced muscle damage. Protective effect of corticosteroids and antioxidants in rabbits. *Acta Orthopaedica Scandinavica* 1996;67(4):393-8.
- ⁴⁰ Son EW, Mo SJ, Rhee DK, Pyo S. Vitamin C blocks TNF-alpha-induced NF-kappaB activation and ICAM-1 expression in human neuroblastoma cells. *Arch Pharm Res.* 2004;27(10):1073-9.
- ⁴¹ Jakeman P, Maxwell S. Effect of antioxidant vitamin supplementation on muscle function after eccentric exercise. *European Journal of Applied Physiology* 1993;67:426-30.
- ⁴² Interactions of vitamin E with free radicals and membranes. Bisby-RH SO: *Free-Radic-Res-Commun.* 1990; 8(4-6): 299-306.
- ⁴³ Maguire-JJ; Wilson-DS; Packer-L. Mitochondrial electron transport-linked tocopheroxyl radical reduction. *J-Biol-Chem.* 1989 Dec 25; 264(36): 21462-5.
- ⁴⁴ Numan-IT; Hassan-MQ; Stohs-SJ. Protective effects of antioxidants against endrin-induced lipid peroxidation, glutathione depletion, and lethality in rats. *Arch-Environ-Contam-Toxicol.* 1990 May-Jun; 19(3): 302-6.
- ⁴⁵ Maellaro-E; Casini-AF; Del-Bello-B; Comporti-M. Lipid peroxidation and antioxidant systems in the liver injury produced by glutathione depleting agents. *Biochem-Pharmacol.* 1990 May 15; 39(10): 1513-21.
- ⁴⁶ Inverse correlation of vitamin E and ischemic heart disease. Gey-KF. *Int-J-Vitam-Nutr-Res-Suppl.* 1989; 30: 224-31.
- ⁴⁷ Yoshida-H; Kajimoto-G. Effect of dietary vitamin E on the toxicity of autoxidized oil to rats. *Ann-Nutr-Metab.* 1989; 33(3): 153-61.
- ⁴⁸ Vitamin E in human low-density lipoprotein. When and how this antioxidant becomes a pro-oxidant. Bowry-VW; Ingold-KU; Stocker-R. *BIOCHEM-J.* 288/2 (341-344) 1992.
- ⁴⁹ Tocopherol mobilization during intensive exercise. Pincemail J; Deby C; Camus G; Pirnay F; Bouchez R; Massaux L; Goutier R. *Eur J Appl Physiol* 1988 57 (2) p189-91.
- ⁵⁰ Kaikkonen J, Kosonen L, Nyssönen K, Porkkala-Sarataho E, Salonen R, Korpela H, Salonen JT. Effect of combined coenzyme Q10 and d-alpha-tocopheryl acetate supplementation on exercise-induced lipid peroxidation and muscular damage: a placebo-controlled double-blind study in marathon runners. *Free Radic Res.* 1998 Jul;29(1):85-92.
- ⁵¹ Sueoka N, Suganuma M, Sueoka E, Okabe S, Matsuyama S, Imai K, Nakachi K, Fujiki H. A new function of green tea: prevention of lifestyle-related diseases. *Ann N Y Acad Sci.* 2001;928:274-80.
- ⁵² Yang F, de Villiers WJ, McClain CJ, Varilek GW. Green tea polyphenols block endotoxin-induced tumor necrosis factor-production and lethality in a murine model. *J Nutr.* 1998;128(12):2334-40.
- ⁵³ Cooper R, Morre DJ, Morre DM. Medicinal benefits of green tea: Part I. Review of noncancer health benefits. *J Altern Complement Med.* 2005;11(3):521-8.
- ⁵⁴ Curtis CL, Harwood JL, Dent CM, Caterson B. Biological basis for the benefit of nutraceutical supplementation in arthritis. *Drug Discov Today.* 2004 15;9(4):165-72.
- ⁵⁵ Ulrich-Vinther M, Maloney MD, Schwarz EM, Rosier R, O'Keefe RJ. Articular cartilage biology. *J Am Acad Orthop Surg.* 2003;11(6):421-30.
- ⁵⁶ Ahmed S, Wang N, Lalonde M, Goldberg VM, Haqqi TM. Green tea polyphenol epigallocatechin-3-gallate (EGCG) differentially inhibits interleukin-1 beta-induced expression of matrix metalloproteinase-1 and -13 in human chondrocytes. *J Pharmacol Exp Ther.* 2004;308(2):767-73.
- ⁵⁷ Haqqi TM. Green tea polyphenol epigallocatechin-3-gallate inhibits the IL-1 beta-induced activity and expression of cyclooxygenase-2 and nitric oxide synthase-2 in human chondrocytes. *Free Radic Biol Med.* 2002;33(8):1097-105.
- ⁵⁸ Adcocks C, Collin P, Buttle DJ. Catechins from green tea (*Camellia sinensis*) inhibit bovine and human cartilage proteoglycan and type II collagen degradation in vitro. *J Nutr.* 2002;132(3):341-6.
- ⁵⁹ Henning SM, Niu Y, Lee NH, Thames GD, Minutti RR, Wang H, Go VL, Heber D. Bioavailability and antioxidant activity of tea flavanols after consumption of green tea, black tea, or a green tea extract supplement. *Am J Clin Nutr.* 2004;80(6):1558-64.
- ⁶⁰ Guardia T, Rotelli AE, Juarez AO, Pelzer LE. Anti-inflammatory properties of plant flavonoids. Effects of rutin, quercetin and hesperidin on adjuvant arthritis in rat. *Farmacol.* 2001;56(9):683-7.

-
- ⁶¹ Teixeira S. Bioflavonoids: proanthocyanidins and quercetin and their potential roles in treating musculoskeletal conditions. *J Orthop Sports Phys Ther* 2002;32(7):357-63.
- ⁶² de Pascual-Teresa S, Johnston KL, DuPont MS, et al. Quercetin metabolites downregulate cyclooxygenase-2 transcription in human lymphocytes ex vivo but not in vivo. *J Nutr* 2004;134(3):552-557.
- ⁶³ Davis JM, Murphy EA, Carmichael MD, Davis B. Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am J Physiol Regul Integr Comp Physiol*. 2009 Apr;296(4):R1071-7.
- ⁶⁴ Duranti G, Ceci R, Patrizio F, Sgrò P, Di Luigi L, Sabatini S, Felici F, Bazzucchi I. Chronic consumption of quercetin reduces erythrocytes oxidative damage: Evaluation at resting and after eccentric exercise in humans. *Nutr Res*. 2018 Feb;50:73-81.
- ⁶⁵ Bazzucchi I, Patrizio F, Ceci R, Duranti G, Sgrò P, Sabatini S, Di Luigi L, Sacchetti M, Felici F. The Effects of Quercetin Supplementation on Eccentric Exercise-Induced Muscle Damage. *Nutrients*. 2019 Jan 21;11(1).
- ⁶⁶ Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: a component of tumeric (*Curcuma longa*). *J Altern Complement Med*. 2003;9(1):161-8.
- ⁶⁷ Satoskar RR, Shah SJ, Shenoy SG. Evaluation of anti-inflammatory property of curcumin (diferuloyl methane) in patients with postoperative inflammation. *Int J Clin Pharmacol Ther Toxicol*. 1986;24(12):651-4.
- ⁶⁸ Alappat L, Awad AB. Curcumin and obesity: evidence and mechanisms. *Nutr Rev*. 2010 Dec;68(12):729-38.
- ⁶⁹ Nicol LM, Rowlands DS, Fazakerly R, Kellett J. Curcumin supplementation likely attenuates delayed onset muscle soreness (DOMS). *Eur J Appl Physiol*. 2015 Aug;115(8):1769-77.
- ⁷⁰ McFarlin BK, Venable AS, Henning AL, Sampson JN, Pennel K, Vingren JL, Hill DW. Reduced inflammatory and muscle damage biomarkers following oral supplementation with bioavailable curcumin. *BBA Clin*. 2016 Feb 18;5:72-8.
- ⁷¹ Davis JM, Murphy EA, Carmichael MD, Zielinski MR, Groschwitz CM, Brown AS, Gangemi JD, Ghaffar A, Mayer EP. Curcumin effects on inflammation and performance recovery following eccentric exercise-induced muscle damage. *Am J Physiol Regul Integr Comp Physiol*. 2007 Jun;292(6):R2168-73.
- ⁷² Delecroix B, Abaïdia AE, Leduc C, Dawson B, Dupont G. Curcumin and Piperine Supplementation and Recovery Following Exercise Induced Muscle Damage: A Randomized Controlled Trial. *J Sports Sci Med*. 2017 Mar 1;16(1):147-153.
- ⁷³ McFarlin BK, Venable AS, Henning AL, Sampson JN, Pennel K, Vingren JL, Hill DW. Reduced inflammatory and muscle damage biomarkers following oral supplementation with bioavailable curcumin. *BBA Clin*. 2016 Feb 18;5:72-8.
- ⁷⁴ Phan TT, See P, Lee ST, Chan SY. Protective effects of curcumin against oxidative damage on skin cells in vitro: its implication for wound healing. *J Trauma*. 2001;51(5):927-31.
- ⁷⁵ Delecroix B, Abaïdia AE, Leduc C, Dawson B, Dupont G. Curcumin and Piperine Supplementation and Recovery Following Exercise Induced Muscle Damage: A Randomized Controlled Trial. *J Sports Sci Med*. 2017 Mar 1;16(1):147-153.
- ⁷⁶ Goozee KG, Shah TM, Sohrabi HR, Rainey-Smith SR, Brown B, Verdile G, Martins RN. Examining the potential clinical value of curcumin in the prevention and diagnosis of Alzheimer's disease. *Br J Nutr*. 2016 Feb 14;115(3):449-65.
- ⁷⁷ Dey A, Bhattacharya R, Mukherjee A, Pandey DK. Natural products against Alzheimer's disease: Pharmaco-therapeutics and biotechnological interventions. *Biotechnol Adv*. 2017 Mar - Apr;35(2):178-216.
- ⁷⁸ Wang Y, Lee KW, Chan FL, Chen S, Leung LK. The red wine polyphenol resveratrol displays bilevel inhibition on aromatase in breast cancer cells. *Toxicol Sci*. 2006 Jul;92(1):71-7.
- ⁷⁹ Wang Y, Leung LK. Pharmacological concentration of resveratrol suppresses aromatase in JEG-3 cells. *Toxicol Lett*. 2007 Sep 28;173(3):175-80.

-
- ⁸⁰ Valenzano DR, Terzibasi E, Genade T, Cattaneo A, Domenici L, Cellerino A. Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-lived vertebrate. *Curr Biol.* 2006 Feb 7;16(3):296-300.
- ⁸¹ Wu R-E, Huang W-C, Liao C-C, et al. Resveratrol protects against physical fatigue and improves exercise performance in mice. *Molecules.* 2013;18(4):4689–702.
- ⁸² Myburgh KH. Polyphenol supplementation: benefits for exercise performance or oxidative stress? *Sports Med.* 2014;44(Suppl 1):57–70.
- ⁸³ Hart N, Sarga L, Csende Z, et al. Resveratrol enhances exercise training responses in rats selectively bred for high running performance. *Food Chem Toxicol.* 2013;61:53–9.
- ⁸⁴ Köbe T1, Witte AV2, Schnelle A1, Tesky VA3, Pantel J3, Schuchardt JP4, Hahn A4, Bohlken J5, Grittner U6, Flöel A7. Impact of Resveratrol on Glucose Control, Hippocampal Structure and Connectivity, and Memory Performance in Patients with Mild Cognitive Impairment. *Front Neurosci.* 2017 Mar 7;11:105. doi: 10.3389/fnins.2017.00105. eCollection 2017.
- ⁸⁵ Kan NW, Ho CS, Chiu YS, Huang WC, Chen PY, Tung YT, Huang CC. Effects of Resveratrol Supplementation and Exercise Training on Exercise Performance in Middle-Aged Mice. *Molecules.* 2016 May 18;21(5). pii: E661.
- ⁸⁶ Cui X1, Jing X2, Wu X1, Yan M1. Protective effect of resveratrol on spermatozoa function in male infertility induced by excess weight and obesity. *Mol Med Rep.* 2016 Nov;14(5):4659-4665.
- ⁸⁷ Mankowski RT1, Anton SD, Buford TW, Leeuwenburgh C. Dietary Antioxidants as Modifiers of Physiologic Adaptations to Exercise. *Med Sci Sports Exerc.* 2015 Sep;47(9):1857-68.
- ⁸⁸ Wang S, Liang X, Yang Q, Fu X, Rogers CJ, Zhu M, et al. Resveratrol induces brownlike adipocyte formation in white fat through activation of AMP-activated protein kinase (AMPK) alpha1. *Int J Obes (Lond)* 2015;39:967–76.
- ⁸⁹ Polley KR1,1, Jenkins N1,1, O'Connor P1,1, McCully K1,1. Influence of exercise training with resveratrol supplementation on skeletal muscle mitochondrial capacity. *Appl Physiol Nutr Metab.* 2016 Jan;41(1):26-32.
- ⁹⁰ Price N L, Gomes A P, Ling A J, Duarte F V, Martin-Montalvo A, North B J, et al. SIRT1 is required for AMPK activation and the beneficial effects of resveratrol on mitochondrial function. *Cell Metab,* 2012. 15(5):675±90.
- ⁹¹ McDermott MM, Leeuwenburgh C, Guralnik JM, Tian L, Sufit R, Zhao L, Criqui MH, Kibbe MR, Stein JH, Effect of Resveratrol on Walking Performance in Older People With Peripheral Artery Disease: The RESTORE Randomized Clinical Trial. Lloyd-Jones D, Anton SD, Polonsky TS, Gao Y, de Cabo R, Ferrucci L. *JAMA Cardiol.* 2017 Apr 12.
- ⁹² Xuan L L, Shi J, Yao C S, Bai J Y, Qu F, Zhang J L, et al. Vam3, a resveratrol dimer, inhibits cigarette smoke-induced cell apoptosis in lungs by improving mitochondrial function[J]. *Acta Pharmacol Sin,* 2014, 35(6):779±791.
- ⁹³ Zhong J, Trevisi L, Urch B, Lin X, Speck M, Coull BA, Liss G, Thompson A, Wu S, Wilson A, Koutrakis P, Silverman F, Gold DR, Baccarelli AA. B-vitamin Supplementation Mitigates Effects of Fine Particles on Cardiac Autonomic Dysfunction and Inflammation: A Pilot Human Intervention Trial. *Sci Rep.* 2017 Apr 3;7:45322.
- ⁹⁴ Xuan L L, Shi J, Yao C S, Bai J Y, Qu F, Zhang J L. Vam3, a resveratrol dimer, inhibits cigarette smoke-induced cell apoptosis in lungs by improving mitochondrial function[J]. *Acta Pharmacol Sin,* 2014, 35(6):779±791.
- ⁹⁵ Nagai K, Suda T. Immunoregulative effects of carnosine and beta-alanine. *J. Physiol. Soc Jap* 1986;48:564-571.
- ⁹⁶ Boldyrev A, Johnson P. Carnosine and related compounds: antioxidant dipeptides. In: P. Johnson and A. Boldyrev, Editors, *Oxidative Stress at Molecular, Cellular and Organ Levels*, Res. Signpost 2002;101-114.
- ⁹⁷ Nagai K, Suda T, Kawasaki K, Mathuura S. Action of carnosine and beta-alanine on wound healing. *Surgery.* 1986;100(5):815-21.
- ⁹⁸ Wang AM, Ma C, Xie ZH, et al. Use of carnosine as a natural anti-senescence drug for human beings. *Biochemistry (Mosc)* 2000;65(7):869-71.

-
- ⁹⁹ Zakharchenko MV, Temnov AV, Kondrashova MN. Effect of carnosine on self-organization of mitochondrial assemblies in rat liver homogenate. *Biochemistry (Mosc)*. 2003;68(9):1002-5.
- ¹⁰⁰ Stvolinsky SL, Dobrota D. Anti-ischemic activity of carnosine. *Biochemistry (Mosc)* 2000;65(7):849-55.
- ¹⁰¹ Baye E, Ukropcova B, Ukropec J, Hipkiss A, Aldini G, de Courten B. Physiological and therapeutic effects of carnosine on cardiometabolic risk and disease. *Amino Acids*. 2016 May;48(5):1131-49.
- ¹⁰² Sale C, Artioli GG, Gualano B, Saunders B, Hobson RM, Harris RC. Carnosine: from exercise performance to health. *Amino Acids*. 2013 Jun;44(6):1477-91.
- ¹⁰³ Baye E, Ukropcova B, Ukropec J, Hipkiss A, Aldini G, de Courten B. Physiological and therapeutic effects of carnosine on cardiometabolic risk and disease. *Amino Acids*. 2016 May;48(5):1131-49.
- ¹⁰⁴ Hipkiss AR. Possible Benefit of Dietary Carnosine towards Depressive Disorders. *Aging Dis*. 2015 Oct 1;6(5):300-3.
- ¹⁰⁵ Lombardi C, Metra M. Carnosine: Potential aid for diabetes and cardiovascular disease. *Obesity (Silver Spring)*. 2016 May;24(5):989.
- ¹⁰⁶ Hipkiss AR, Baye E, de Courten B. Carnosine and the processes of ageing. *Maturitas*. 2016 Nov;93:28-33.
- ¹⁰⁷ Baye E, Ukropcova B, Ukropec J, Hipkiss A, Aldini G, de Courten B. Physiological and therapeutic effects of carnosine on cardiometabolic risk and disease. *Amino Acids*. 2016 May;48(5):1131-49.
- ¹⁰⁸ Bellia F, Vecchio G, Cuzzocrea S, Calabrese V, Rizzarelli E. Neuroprotective features of carnosine in oxidative driven diseases. *Mol Aspects Med*. 2011 Aug;32(4-6):258-66.
- ¹⁰⁹ Dutka, T.L., Lambole, C.R., McKenna, M.J., Murphy, R.M., and Lamb, G.D. 2012. Effects of carnosine on contractile apparatus Ca(2)(+) sensitivity and sarcoplasmic reticulum Ca(2)(+) release in human skeletal muscle fibers. *J. Appl. Physiol*. 112(5): 728–736.
- ¹¹⁰ Szczesniak D, Budzen S, Kopec W, Rymaszewska J. Anserine and carnosine supplementation in the elderly: Effects on cognitive functioning and physical capacity. *Arch Gerontol Geriatr*. 2014 Sep-Oct;59(2):485-90.
- ¹¹¹ Liu Y, Cotillard A, Vazier C, Bastard J, Fellahi S, Stévant M, Allatif O, Langlois C, Bieuvelet S, Brochot A, Guilbot A, Clément K, Rizkalla S (2015) A dietary supplement containing cinnamon, chromium and carnosine decreases fasting plasma glucose and increases lean mass in overweight or obese pre-diabetic subjects: a randomized, placebo-controlled trial. *PLoS One* 10(12)
- ¹¹² Anderson RA. Chromium and polyphenols from cinnamon improve insulin sensitivity. *Proc Nutr Soc*. 2008 Feb;67(1):48-53.
- ¹¹³ Mortensen SA. Perspectives on therapy of cardiovascular diseases with coenzyme Q10 (ubiquinone). [Review] *Clinical Investigator* 1993;71(8 Suppl):S116-23.
- ¹¹⁴ Beyer RE. An analysis of the role of coenzyme Q in free radical generation and as an antioxidant. *Biochemistry & Cell Biology* 1992;70(6):390-403.
- ¹¹⁵ Chis BA, Chis AF, Muresan A, Fodor D. Q10 Coenzyme Supplementation can Improve Oxidative Stress Response to Exercise in Metabolic Syndrome in Rats. *Int J Vitam Nutr Res*. 2019 Mar 19:1-9.
- ¹¹⁶ Belviranlı M, Okudan N. Effect of Coenzyme Q10 Alone and in Combination with Exercise Training on Oxidative Stress Biomarkers in Rats. *Int J Vitam Nutr Res*. 2019 Apr 30:1-11.
- ¹¹⁷ Sarmiento A, Diaz-Castro J, Pulido-Moran M, Moreno-Fernandez J, Kajarabille N, Chiroso I, Guisado IM, Javier Chiroso L, Guisado R, Ochoa JJ. Short-term ubiquinol supplementation reduces oxidative stress associated with strenuous exercise in healthy adults: A randomized trial. *Biofactors*. 2016 Nov 12;42(6):612-622.
- ¹¹⁸ Orlando P, Silvestri S, Galeazzi R, Antonicelli R, Marcheggiani F, Cirilli I, Bacchetti T, Tiano L. Effect of ubiquinol supplementation on biochemical and oxidative stress indexes after intense exercise in young athletes. *Redox Rep*. 2018 Dec;23(1):136-145.
- ¹¹⁹ Borisova IG, Seifulla RD, Zhuravlev AI. [Action of antioxidants on physical work capacity and lipid peroxidation in the body]. *Farmakol Toksikol* 1989;52(4):89-92.
- ¹²⁰ Lemke M, Frei B, Ames BN, Faden AI. Decreases in tissue levels of ubiquinol 9 and 10, ascorbate and alpha tocopherol following spinal cord impact trauma in rats. *Neurosci Lett* 1990;108(1-2):201-6.

-
- ¹²¹ Bendich A. Symposium conclusions: biological actions of carotenoids. *J Nutr*; 1989 Jan; 119(1); P 135-6.
- ¹²² Mayne ST, Janerich DT, Greenwald P, Chorost S, Tucci C, Zaman MB, Melamed MR, Kiely M, McKneally MF. Dietary beta carotene and lung cancer risk in U.S. nonsmokers. *Journal of the National Cancer Institute*. 86(1):33-8, 1994 Jan 5.
- ¹²³ Preferential inhibition of LDL oxidation by the all-trans isomer of beta -carotene in comparison with 9-cis beta -carotene Lavy-A; Ben-Amotz-A; Aviram-M. *EUR-J-CLIN-CHEM-CLIN-BIOCHEM*. 31/2 (83-90) 1993.
- ¹²⁴ Dawson Jr R, Biasseti M, Messina S, Dominy J. The cytoprotective role of taurine in exercise-induced muscle injury. *Amino Acids* 2002;22(4):309-24.
- ¹²⁵ Manabe S, Kurroda I, Okada K, Morishima M, Okamoto M, Harada N, Takahashi A, Sakai K, Nakaya Y. Decreased blood levels of lactic acid and urinary excretion of 3-methylhistidine after exercise by chronic taurine treatment in rats. *J Nutr Sci Vitaminol (Tokyo)*. 2003 Dec;49(6):375-80.
- ¹²⁶ Zhang M, Izumi I, Kagamimori S, Sokejima S, Yamagami T, Liu Z, Qi B. Role of taurine supplementation to prevent exercise-induced oxidative stress in healthy young men. *Amino Acids*. 2004 Mar;26(2):203-7.
- ¹²⁷ Abbasoglu L, Kalaz EB, Soluk-Tekkesin M, Olgaç V, Dogru-Abbasoglu S, Uysal M. Beneficial effects of taurine and carnosine in experimental ischemia/reperfusion injury in testis. *Pediatr Surg Int*. 2012 Nov;28(11):1125-31.
- ¹²⁸ Fraser SF, Li JL, Carey MF, Wang XN, Sangkabutra T, Sostaric S, Selig SE, Kjeldsen K, McKenna MJ. Fatigue depresses maximal in vitro skeletal muscle Na(+)-K(+)-ATPase activity in untrained and trained individuals. *J Appl Physiol* 2002 Nov;93(5):1650-9.
- ¹²⁹ Li JL, Wang XN, Fraser SF, Carey MF, Wrigley TV, McKenna MJ. Effects of fatigue and training on sarcoplasmic reticulum Ca(2+) regulation in human skeletal muscle. *J Appl Physiol* 2002;92(3):912-22.
- Abbasoglu L, Kalaz EB, Soluk-Tekkesin M, Olgaç V, Dogru-Abbasoglu S, Uysal M. Beneficial effects of taurine and carnosine in experimental ischemia/reperfusion injury in testis. *Pediatr Surg Int*. 2012 Nov;28(11):1125-31. ¹³⁰ Yatabe Y, Miyakawa S, Miyazaki T, Matsuzaki Y, Ochiai N. Effects of taurine administration in rat skeletal muscles on exercise. *J Orthop Sci*. 2003;8(3):415-9.
- ¹³¹ Bast A, Haenen GR. Lipoic acid: a multifunctional antioxidant. *Biofactors*. 2003;17(1-4):207-13.
- ¹³² Packer L, Witt EH, Tritschler HJ. Alpha-lipoic acid as a biological antioxidant. *Free Radic Biol Med*. 1995;19:227-250.
- ¹³³ Jones W, Li X, Qu ZC, et al. Uptake, recycling, and antioxidant actions of alpha-lipoic acid in endothelial cells. *Free Radic Biol Med* 2002;33:83-93.
- ¹³⁴ Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Radic Biol Med* 1997;22(1-2):359-78.
- ¹³⁵ Podda M, Tritschler HJ, Ulrich H, et al. Alpha-lipoic acid supplementation prevents symptoms of vitamin E deficiency. *Biochem Biophys Res Commun*. 1994;204:98-104.
- ¹³⁶ Patrick L. Mercury toxicity and antioxidants: Part 1: role of glutathione and alpha-lipoic acid in the treatment of mercury toxicity. *Altern Med Rev*. 2002;7(6):456-71.
- ¹³⁷ Haghghatdoost F, Hariri M. The effect of alpha-lipoic acid on inflammatory mediators: a systematic review and meta-analysis on randomized clinical trials. *Eur J Pharmacol*. 2019 Apr 15;849:115-123.
- ¹³⁸ Maroon JC, Bost JW. Omega-3 fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain. *Surg Neurol*. 2006;65(4):326-31.
- ¹³⁹ Packer L. Alpha lipoic acid: a metabolic antioxidant which regulates NF- kappaB signal transduction and protects against oxidative injury. *Drug Metab Rev* 1998;30:245-75.
- ¹⁴⁰ Lee HA, Hughes DA. Alpha-lipoic acid modulates NF-kappaB activity in human monocytic cells by direct interaction with DNA. *Exp Gerontol*. 2002;37(2-3):401-10.
- ¹⁴¹ Mousavi SM, Shab-Bidar S, Kord-Varkaneh H, Khorshidi M, Djafarian K. Effect of alpha-lipoic acid supplementation on lipid profile: A systematic review and meta-analysis of controlled clinical trials. *Nutrition*. 2019 Mar;59:121-130.

-
- ¹⁴² Tromba L, Perla FM, Carbotta G, Chiesa C, Pacifico L. Effect of Alpha-Lipoic Acid Supplementation on Endothelial Function and Cardiovascular Risk Factors in Overweight/Obese Youths: A Double-Blind, Placebo-Controlled Randomized Trial. *Nutrients*. 2019 Feb 12;11(2). pii: E375.
- ¹⁴³ Faust A, Burkart V, Ulrich H, Weischer CH, Kolb H. Effect of lipoic acid on cyclophosphamide-induced diabetes and insulinitis in non-obese diabetic mice. *Int J Immunopharmacol*. 1994;16(1):61-6.
- ¹⁴⁴ Burkart V, Koike T, Brenner HH, Imai Y, Kolb H. Dihydrolipoic acid protects pancreatic islet cells from inflammatory attack. *Agents Actions*. 1993;38(1-2):60-5.
- ¹⁴⁵ Lateef H, Aslam MN, Stevens MJ, Varani J. Pretreatment of diabetic rats with lipoic acid improves healing of subsequently-induced abrasion wounds. *Arch Dermatol Res* 2005;297(2):75-83.
- ¹⁴⁶ Thirunavukkarasu V, Nandhini AT, Anuradha CV. Fructose diet-induced skin collagen abnormalities are prevented by lipoic acid. *Exp Diabetes Res*. 2004;5(4):237-44.
- ¹⁴⁷ Li N, Yan W, Hu X, Huang Y, Wang F, Zhang W, Wang Q, Wang X, Sun K. Effects of oral alpha-lipoic acid administration on body weight in overweight or obese subjects: a crossover randomized, double-blind, placebo-controlled trial. *Clin Endocrinol (Oxf)*. 2017 May;86(5):680-687.
- ¹⁴⁸ Arivazhagan P, Ramanathan K, Panneerselvam C. Effect of DL-alpha-lipoic acid on mitochondrial enzymes in aged rats. *Chem Biol Interact*. 2001 Nov 28;138(2):189-98.
- ¹⁴⁹ Palaniappan AR, Dai A. Mitochondrial ageing and the beneficial role of alpha-lipoic acid. *Neurochem Res*. 2007 Sep;32(9):1552-8.
- ¹⁵⁰ Margaritis I, Rousseau AS, Hininger I, Palazzetti S, Arnaud J, Roussel AM. Increase in selenium requirements with physical activity loads in well-trained athletes is not linear. *Biofactors*. 2005;23(1):45-55. Links
- ¹⁵¹ Lukaski HC: Low dietary zinc decreases erythrocyte carbonic anhydrase activities and impairs cardiorespiratory function in men during exercise. *Am J Clin Nutr* 2005; 81: 1045-1051.
- ¹⁵² Prasad AS. Zinc deficiency in women, infants and children. *Journal of the American College of Nutrition* 1996;15(2):113-20.
- ¹⁵³ Cordova A, Alvarez-Mon M. Behaviour of zinc in physical exercise: a special reference to immunity and fatigue. *Neuroscience & Biobehavioral Reviews* 1995;19(3):439-45.
- ¹⁵⁴ Kieffer F. (Trace elements: their importance for health and physical performance.) *Deutsche Zeitschrift fuer Sportmedizin* 1986;37(4):118-123.
- ¹⁵⁵ Oteiza PI, Olin KL, Fraga CG, Keen CL. Zinc deficiency causes oxidative damage to proteins, lipids and DNA in rat testes. *J Nutr* 1995;125(4):823-9.
- ¹⁵⁶ Hsu JM. Zinc deficiency and alterations of free amino acid levels in plasma, urine and skin extract. *Progress in Clinical & Biological Research* 1977;14:73-86.
- ¹⁵⁷ Dorup I, Flyvbjerg A, Everts ME, Clausen T. Role of insulin-like growth factor-1 and growth hormone in growth inhibition induced by magnesium and zinc deficiencies. *British Journal of Nutrition* 1991;66(3):505-21.
- ¹⁵⁸ Ghavami-Maibodi SZ, Collipp PJ, Castro-Magana M, Stewart C and Chen SY. Effect of oral zinc supplements on growth, hormonal levels and zinc in healthy short children. *Ann Nutr Metab* 1983;273:214-219.
- ¹⁵⁹ Hartoma TR, Nahoul K, Netter A. Zinc, plasma androgens and male sterility. *Lancet* 1977;2:1125-1126.
- ¹⁶⁰ Hunt CD, Johnson PE, Herbel J, Mullen LK. Effects of dietary zinc depletion on seminal volume of zinc loss, serum testosterone concentrations and sperm morphology in young men. *Am J Clin Nutr* 1992;56(1):148-157.
- ¹⁶¹ Kara E, Gunay M, Cicioglu I, Ozal M, Kilic M, Mogulkoc R, Baltaci AK. Effect of zinc supplementation on antioxidant activity in young wrestlers. *Biol Trace Elem Res*. 2010 Apr;134(1):55-63.
- ¹⁶² Marreiro DN, Geloneze B, Tambascia MA, Lerario AC, Halpern A, Cozzolino SM. [Participation of zinc in insulin resistance] *Arq Bras Endocrinol Metabol*. 2004;48(2):234-9.
- ¹⁶³ Lelovics Z. Relation between calcium and magnesium intake and obesity. *Asia Pac J Clin Nutr*. 2004;13(Suppl):S144.

-
- ¹⁶⁴ Rayssiguier Y, Mazur A. R [Magnesium and inflammation: lessons from animal models.] Clin Calcium. 2005;15(2):245-248.
- ¹⁶⁵ Maier JA, Malpuech-Brugere C, Zimowska W, Rayssiguier Y, Mazur A. Low magnesium promotes endothelial cell dysfunction: implications for atherosclerosis, inflammation and thrombosis. Biochim Biophys Acta. 2004 May 24;1689(1):13-21.
- ¹⁶⁶ Seelig MS. Consequences of magnesium deficiency on the enhancement of stress reactions; preventive and therapeutic implications (a review). J Am Coll Nutr. 1994 Oct;13(5):429-46.
- ¹⁶⁷ Nielsen FH, Lukaski HC. Update on the relationship between magnesium and exercise. Magnes Res. 2006 Sep;19(3):180-9.
- ¹⁶⁸ Setaro L, Santos-Silva PR, Nakano EY, Sales CH, Nunes N, Greve JM, Colli C. Magnesium status and the physical performance of volleyball players: effects of magnesium supplementation. J Sports Sci. 2014;32(5):438-45.
- ¹⁶⁹ Santos DA, Matias CN, Monteiro CP, Silva AM, Rocha PM, Minderico CS, Bettencourt Sardinha L, Laires MJ. Magnesium intake is associated with strength performance in elite basketball, handball and volleyball players. Magnes Res. 2011 Dec;24(4):215-9.
- ¹⁷⁰ Garrison SR, Allan GM, Sekhon RK, Musini VM, Khan KM. Magnesium for skeletal muscle cramps. Cochrane Database Syst Rev. 2012 Sep 12;9:CD009402.
- ¹⁷¹ Maggio M, De Vita F, Lauretani F, Nouvenne A, Meschi T, Ticinesi A, Dominguez LJ, Barbagallo M, Dall'aglio E, Ceda GP. The Interplay between Magnesium and Testosterone in Modulating Physical Function in Men. Int J Endocrinol. 2014;2014:525249.
- ¹⁷² Maggio M, Ceda GP, Lauretani F, Cattabiani C, Avantageggiato E, Morganti S, Ablondi F, Bandinelli S, Dominguez LJ, Barbagallo M, Paolisso G, Semba RD, Ferrucci L. Magnesium and anabolic hormones in older men. Int J Androl. 2011 Dec;34(6 Pt 2):e594-600.
- ¹⁷³ Koehler K, Parr MK, Geyer H, Mester J, Schänzer W. Serum testosterone and urinary excretion of steroid hormone metabolites after administration of a high-dose zinc supplement. Eur J Clin Nutr. 2009 Jan;63(1):65-70.
- ¹⁷⁴ K. Tajima et al., antioxidant activity of Gamma Oryzanol. Biochem. Biophys. Res. Commun. 115, 1002 (1983).
- ¹⁷⁵ Inhibitory effect of taurine on HOCl- and NH₂Cl-induced degradation of hyaluronic acid Lee-C-S; Lee-K-Y; Lee-K-S. KOREAN-J-PHARMACOL. 28/2 (201-212) 1992.
- ¹⁷⁶ Histidine dipeptide levels in ageing and hypertensive rat skeletal and cardiac muscles. Johnson-P; Hammer-JL. COMP-BIOCHEM-PHYSIOL-B-COMP-BIOCHEM. 103/4 (981-984) 1992.
- ¹⁷⁷ Zeynep Tuzcu, Cemal Orhan, Nurhan Sahin, Kazim Sahin, Cinnamon Extract Inhibit Lipids and Inflammation by Modulation of Transcription Factors: in vivo Model American Heart Association's Arteriosclerosis, Thrombosis and Vascular Biology | Peripheral Vascular Disease 2017 Scientific Sessions; 545.
- ¹⁷⁸ Frei-B; Kim-MC; Ames-BN. Ubiquinol-10 is an effective lipid-soluble antioxidant at physiological concentrations. Proc-Natl-Acad-Sci-U-S-A. 1990 Jun; 87(12): 4879-83.
- ¹⁷⁹ Lemke-M; Frei-B; Ames-BN; Faden-AI. Decreases in tissue levels of ubiquinol-9 and -10, ascorbate and alpha-tocopherol following spinal cord impact trauma in rats. Neurosci-Lett. 1990 Jan 1; 108(1-2): 201-6.
- ¹⁸⁰ Sato-K; Niki-E; Shimasaki-H. Free radical-mediated chain oxidation of low density lipoprotein and its synergistic inhibition by vitamin E and vitamin C. Arch-Biochem-Biophys. 1990 Jun; 279(2): 402-5.
- ¹⁸¹ van-den-Berg-JJ; Kuypers-FA; Roelofsen-B; Op-den-Kamp-JA. The cooperative action of vitamins E and C in the protection against peroxidation of parinaric acid in human erythrocyte membranes. Chem-Phys-Lipids. 1990 Mar; 53(4): 309-20.
- ¹⁸² Relationship between free radicals and vitamins: an overview. Sies-H SO: Int-J-Vitam-Nutr-Res-Suppl. 1989; 30: 215-23.
- ¹⁸³ Antioxidant functions of vitamins. Vitamins E and C, beta-carotene, and other carotenoids. Sies H; Stahl W; Sundquist AR. Ann N Y Acad Sci (UNITED STATES) Sep 30 1992, 669 p7-20.

-
- 184 Vatassery-GT; Smith-WE; Quach-HT. Ascorbic acid, glutathione and synthetic antioxidants prevent the oxidation of vitamin E in platelets. *Lipids*. 1989 Dec; 24(12): 1043-7.
- 185 Sies H. Oxidative stress: from basic research to clinical application, *Am J Med* 1991; 91:31S–38S.
- 186 Pinsky-C; Bose-R. Pyridine and other coal tar constituents as free radical-generating environmental neurotoxicants. *Mol Cell Biochem* 1988. vol. 84(2), 217-222.
- 187 Fox DL. Air pollution. *Anal Chem*. 1999 Jun 15;71(12):109R-119R.
- 188 Sydbom A, Blomberg A, Parnia S, Stenfors N, Sandstrom T, Dahlen SE. Health effects of diesel exhaust emissions. *Eur Respir J*. 2001 Apr;17(4):733-46.
- 189 Fridovich I. Superoxide dismutases. *Annu Rev Biochem* 1975; 44:147-59.
- 190 Mylonas C, Kouretas D. Lipid peroxidation and tissue damage. *In Vivo*. 1999 May-Jun;13(3):295-309.
- 191 Pryor WA. The formation of free radicals and the consequences of their reactions in vivo. *Photochem Photobiol* 1978; 28:787-801.
- 192 Jessup-W; Rankin-SM; De-Whalley-CV; Hoult-JR; Scott-J; Leake-DS. Alpha-tocopherol consumption during low-density-lipoprotein oxidation. *Biochem-J*. 1990 Jan 15; 265(2): 399-405.
- 193 Endogenous DNA damage as related to cancer and aging. Ames BN Department of Biochemistry, University of California, Berkeley 94720. *Mutat Res* 1989, 214 (1) p41-6.
- 194 Leibovitz BE, Siegel BV. Aspects of free radical reactions in biological systems: aging. *J Gerontol* 1980; 35:45-56.
- 195 Rikans LE, Hornbrook KR. Lipid peroxidation, antioxidant protection and aging. *Biochim Biophys Acta*. 1997 Dec 31;1362(2-3):116-27.
- 196 Ames BN, Cathcart R, Schwiers E, Hochstein P. Uric acid provides an antioxidant defense against oxidant- and radical-caused aging and cancer: a hypothesis. *Proc Natl Acad Sci USA* 1981; 78:6858-62.
- 197 Radi R. Biological antioxidant defenses. *Toxicol Ind Health*. 1993 Jan-Apr;9(1-2):53-62.
- 198 Harman D. Aging: a theory based on free radical and radiation chemistry. *J Gerontol* 1956; 11:298-300.
- 199 Harman D. Free radical theory of aging: history. *EXS*. 1992;62:1-10.
- 200 Harman D. Free-radical theory of aging. Increasing the functional life span. *Ann N Y Acad Sci*. 1994 Jun 30;717:1-15.
- 201 Packer-L; Landvik-S. I. Vitamin E: introduction to biochemistry and health benefits. *Ann-N-Y-Acad-Sci*. 1989; 570: 1-6.
- 202 Klotz LO, K.D. Kröncke, D.P. Buchczyk and H. Sies, Role of copper, zinc, selenium and tellurium in the cellular defense against oxidative and nitrosative stress, *J. Nutr*. 2003; 133:1448S–1451S.
- 203 Miranda-Vilela AL. Exercise, injuries and athlete performance. In: Bastos JH and Silva AC (eds) *Athlete Performance and Injuries*. 2012; Nova Science Publishers, New York, pp 1-50.
- 204 Miranda-Vilela AL, Ribeiro IF, Grisolia CK. Association between interleukin 6 -174 G/C promoter gene polymorphism and runners' responses to the dietary ingestion of antioxidant supplementation based on pequi (*Caryocar brasiliense* Camb.) oil: a before-after study. *Genet Mol Biol*. 2016 Oct-Dec;39(4):554-566.
- 205 Vincent HK, Bourguignon CM, Vincent KR, Weltman AL, Bryant M, Taylor AG. Antioxidant supplementation lowers exercise-induced oxidative stress in young overweight adults. *Obesity (Silver Spring)*. 2006 Dec;14(12):2224-35.
- 206 Mankowski RT1, Anton SD, Buford TW, Leeuwenburgh C. Dietary Antioxidants as Modifiers of Physiologic Adaptations to Exercise. *Med Sci Sports Exerc*. 2015 Sep;47(9):1857-68.
- 207 Aguiló A, Tauler P, Sureda A, Cases N, Tur J, Pons A. Antioxidant diet supplementation enhances aerobic performance in amateur sportsmen. *J Sports Sci*. 2007 Sep;25(11):1203-10.
- 208 Sangha O, Stucki G. Vitamin E in the treatment of rheumatic diseases. *Zeitschrift für Rheumatologie* 1998;57(4):207-214.
- 209 Satchek JM, Blumberg JB. Role of vitamin E and oxidative stress in exercise. *Nutrition* 2001;(10):809-14.

-
- ²¹⁰ Zafarullah M, Li WQ, Sylvester J, Ahmad M. Molecular mechanisms of N-acetylcysteine actions. *Cell Mol Life Sci* 2003;60(1):6-20.
- ²¹¹ Andriamanalijaona R, Kypriotou M, Bauge C, Renard E, Legendre F, Raoudi M, Boumediene K, Gatto H, Monginoux P, Pujol JP. Comparative effects of 2 antioxidants, selenomethionine and epigallocatechin-gallate, on catabolic and anabolic gene expression of articular chondrocytes. *J Rheumatol.* 2005;32(10):1958-67.
- ²¹² Makker K, Agarwal A, Sharma R. Oxidative stress & male infertility. *Indian J Med Res.* 2009 Apr;129(4):357-67.
- ²¹³ Agarwal A, Gupta S, Sharma RK. Role of oxidative stress in female reproduction. *Reprod Biol Endocrinol.* 2005 Jul 14;3:28.
- ²¹⁴ Davis JM, Murphy EA, Carmichael MD, Davis B. Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am J Physiol Regul Integr Comp Physiol.* 2009 Apr;296(4):R1071-7.
- ²¹⁵ Kinnunen S, Hyyppä S, Oksala N, Laaksonen DE, Hannila ML, Sen CK, Atalay M. alpha-Lipoic acid supplementation enhances heat shock protein production and decreases post exercise lactic acid concentrations in exercised standardbred trotters. *Res Vet Sci.* 2009 Nov;6(4):369-95.
- ²¹⁶ Wang XM; Xie ZF. [A clinical study of the effect of wuzi yanzong solution in retarding aging process] *Chung Kuo Chung Hsi I Chieh Ho Tsa Chih (CHINA)* Jan 1992, 12 (1) p23-5.
- ²¹⁷ Borek-C; Maltoni-C; Selikoff-IJ eds. Ozone carcinogenesis in vitro and its co-carcinogenesis with radiation: Discussion paper. *Living In A Chemical World: Occupational And Environmental Significance Of Industrial Carcinogens.* Ann. N.y. Acad. Sci. 1988. pp. 106-110.
- ²¹⁸ Zhang L. Blot WJ. You WC. Chang YS. Liu XQ. Kneller RW. Zhao L. Liu WD. Li JY. Jin ML. et al. Serum micronutrients in relation to pre-cancerous gastric lesions. *International Journal of Cancer.* 56(5):650-4, 1994.
- ²¹⁹ Preventive potential of antioxidative vitamins and carotenoids on cancer. Stahelin-HB; Gey-F; Brubacher-G. *Int-J-Vitam-Nutr-Res-Suppl.* 1989; 30: 232-41.
- ²²⁰ [Relationship between atherosclerosis and aging process]. Kuzuya-F. *Rinsho-Byori.* 1989 Sep; 37(9): 968-74.
- ²²¹ Vitamin E content and low density lipoprotein oxidizability induced by free radicals. Babiy-AV; Gebicki-JM; Sullivan-DR. *Atherosclerosis.* 1990 Apr; 81(3): 175-82.
- ²²² Tappel AL, Fletcher B, Deamer D. Effect of antioxidants and nutrients on lipid peroxidation fluorescent products and aging parameters in the mouse. *J Gerontol* 1973; 28:415-24.
- ²²³ Sukolinskii-VN; Morozkina-TS. [Prevention of postoperative complications in patients with stomach cancer using an antioxidant complex]. *Vopr-Onkol.* 1989; 35(10): 1242-5.
- ²²⁴ A multivitamin infusion prevents lipid peroxidation and improves transplantation performance Rahl-H; Khoschsorur-G; Colombo-T; Petritsch-P; Rauchenwald-M; Koltringer-P; Tatzber-F; Esterbauer-H *KIDNEY-INT.* 43/4 (912-917) 1993.
- ²²⁵ Goode HF. Webster NR. Howdle PD. Leek JP. Lodge JP. Sadek SA. Walker BE. Reperfusion injury, antioxidants and hemodynamics during orthotopic liver transplantation. *Hepatology.* 19(2):354-9, 1994 Feb.
- ²²⁶ Ek B. Hallberg C. Sjogren KG. Hjalmarsen A. ASTRA HASSLE AB, Molndal, Sweden. The Reoxygenation-induced cell damage of isolated neonatal rat ventricular myocytes can be reduced by chain-breaking antioxidants. *Free Radical Biology & Medicine.* 16(1):117-21, 1994 Jan.
- ²²⁷ Vitamin E intake and risk of cataracts in humans. Robertson-JM; Donner-AP; Trevithick-JR. *Ann-N-Y-Acad-Sci.* 1989; 570: 372-82.
- ²²⁸ Gerster H. Antioxidant vitamins in cataract prevention. *Z Ernahrungswiss;* 1989 28(1); P 56-75.
- ²²⁹ The role of antioxidant nutrients in preventing hyperbaric oxygen damage to the retina. Stone WL; Henderson RA; Howard GH Jr; Hollis AL; Payne PH; Scott RL. *Free Radic Biol Med* 1989, 6 (5) p505-12.
- ²³⁰ Cataract: Relationships between nutrition and oxidation Taylor-A *J-AM-COLL-NUTR.* 12/2 (138-146) 1993.

-
- ²³¹ Heliovaara M. Knekt P. Aho K. Aaran RK. Alfthan G. Aromaa A. Serum antioxidants and risk of rheumatoid arthritis. *Source Annals of the Rheumatic Diseases*. 53(1):51-3, 1994.
- ²³² Paolisso G. Gambardella A. Galzerano D. Varricchio M. D'Onofrio F. Antioxidants in adipose tissue and risk of myocardial infarction [letter]. *Source Lancet*. 343(8897):596, 1994 Mar 5.
- ²³³ Weisel-RD; Mickle-DA; Finkle-CD; Tumiaty-LC; Madonik-MM; Ivanov-J; Burton-GW; Ingold-KU. Myocardial free-radical injury after cardioplegia. *Circulation*. 1989 Nov; 80(5 Pt 2): III14-8.
- ²³⁴ Riemersma-RA; Wood-DA; Macintyre-CC; Elton-R; Gey-KF; Oliver-MF. Low plasma vitamins E and C. Increased risk of angina in Scottish men. *Ann-N-Y-Acad-Sci*. 1989; 570: 291-5.
- ²³⁵ Reactive oxygen in skeletal muscle. I. Intracellular oxidant kinetics and fatigue in vitro. Reid-MB; Haack-KE; Franchek-KM; Valberg-PA; Kobzik-L; West-MS. *J-APPL-PHYSIOL*. 73/5 (1797-1804) 1992.
- ²³⁶ Gohil K; Rothfuss L; Lang J; Packer L. Effect of exercise training on tissue vitamin E and ubiquinone content. *J Appl Physiol* 1987 63 (4) p1638-41.
- ²³⁷ Vats P, Singh VK, Singh SN, Singh SB. Glutathione metabolism under high-altitude stress and effect of antioxidant supplementation. *Aviat Space Environ Med*. 2008 Dec;79(12):1106-11.
- ²³⁸ Jayalakshmi K, Singh SB, Kalpana B, Sairam M, Muthuraju S, Ilavazhagan G. N-acetyl cysteine supplementation prevents impairment of spatial working memory functions in rats following exposure to hypobaric hypoxia. *Physiol Behav*. 2007 Nov 23;92(4):643-50.
- ²³⁹ Antioxidant enzyme response to exercise and aging. AU: Li-Li-Ji. *MED-SCI-SPORTS-EXERC*. 25/2 (225-231) 1993.
- ²⁴⁰ Skeletal muscle oxidative capacity, antioxidant enzymes, and exercise training. Laughlin MH; Simpson T; Sexton WL; Brown OR; Smith JK; Korthuis RJ. *J Appl Physiol* Jun 1990, 68 (6) p2337-43.
- ²⁴¹ Antioxidants and the influence of free radical damage to proteins on proteolysis in and around mammalian cells. Naish-Byfield-S; Dean-RT. *Revis-Biol-Celular*. 1989; 21: 361-75.
- ²⁴² Robertson JD, Maughan RJ, Duthie GG, Morrice PC. Blood antioxidant status and erythrocyte lipid peroxidation following distance running. *Arch Biochem Biophys* 1990;282;78-83.
- ²⁴³ Biochemical mechanisms for oxygen free radical formation during exercise. Sjodin B; Hellsten Westing Y; Apple FS. *Sports Med* Oct 1990, 10 (4) p236-54.
- ²⁴⁴ Asayama-K; Kato-K. Oxidative muscular injury and its relevance to hyperthyroidism. *Free-Radic-Biol-Med*. 1990; 8(3): 293-303.
- ²⁴⁵ Asayama-K; Dobashi-K; Hayashibe-H; Kato-K. Vitamin E protects against thyroxine-induced acceleration of lipid peroxidation in cardiac and skeletal muscles in rats. *J-Nutr-Sci-Vitaminol-Tokyo*. 1989 Oct; 35(5): 407-18.
- ²⁴⁶ Brown fat thermogenesis and exercise: two examples of physiological oxidative stress? Barja de Quiroga G. *Free Radic Biol Med* 1992, 13 (4) p325-40.
- ²⁴⁷ Barclay JK; Hansel M. Free radicals may contribute to oxidative skeletal muscle fatigue. *Can J Physiol Pharmacol* Feb 1991, 69 (2) p279-84.
- ²⁴⁸ Novelli GP, Bracciotti G, Falsini S. Spn-trappers and vitamin E prolong endurance to muscle fatigue in mice. *Free Rad Biol Med* 8 1990 pg 236-242.
- ²⁴⁹ Reid, M. B.; Stokic, D. S.; Koch, S. M.; Khawli, F. A.; Leis, A. A. N-acetylcysteine inhibits muscle fatigue in humans. *J. Clin. Invest*. 94: 2468-2474; 1994.
- ²⁵⁰ Medved, I.; Brown, M. J.; Bjorksten, A. R.; Murphy, K. T.; Petersen, A. C.; Sostaric, S.; Gong, X.; McKenna, M. J. N-acetylcysteine enhances muscle cysteine and glutathione availability and attenuates fatigue during prolonged exercise in endurance-trained individuals. *J. Appl. Physiol*. 97: 1477-1485; 2004.
- ²⁵¹ Matuszczak, Y.; Farid, M.; Jones, J.; Lansdowne, S.; Smith, M. A.; Taylor, A. A.; Reid, M. B. Effects of N-acetylcysteine on glutathione oxidation and fatigue during handgrip exercise. *Muscle Nerve* 32: 633-638; 2005.
- ²⁵² Vinciguerra G, Belcaro G, Bonanni E, Cesarone MR, Rotondi V, Ledda A, Hosoi M, Dugall M, Cacchio M, Cornelli U. Evaluation of the effects of supplementation with Pycnogenol® on fitness in

-
- normal subjects with the Army Physical Fitness Test and in performances of athletes in the 100-minute triathlon. *J Sports Med Phys Fitness*. 2013 Dec;53(6):644-54.
- ²⁵³ Janero-DR; Burghardt-B. Oxidative injury to myocardial membrane: direct modulation by endogenous alpha-tocopherol. *J-Mol-Cell-Cardiol*. 1989 Nov; 21(11): 1111-24.
- ²⁵⁴ Singh-V-N. A current perspective on nutrition and exercise. *J-Nutr*. 1992 Mar. 122(3 Suppl). P 760-5.
- ²⁵⁵ Dragan-I. Dinu-V. Cristea-E. Mohora-N. Ploesteanu-E. Stroescu-V. Studies regarding the effects of an antioxidant compound in top athletes. *Rev-Roum-Physiol*. 1991 Jul-Dec. 28(3-4). P 105-8.
- ²⁵⁶ Effects of an antioxidant vitamin mixture on lipid peroxidation at rest and postexercise Kanter-MM; Nolte-LA; Holloszy-JO *J-APPL-PHYSIOL*. 74/2 (965-969) 1993.
- ²⁵⁷ Singh-V-N. A current perspective on nutrition and exercise. *J-Nutr*. 1992 Mar. 122(3 Suppl). P 760-5. Nutrition has always been a subject of great interest to athletes.
- ²⁵⁸ Kinuta-Y. [Lipid peroxidation and changes in xanthine oxidase in cerebral ischemia]. *Nippon-Geka-Hokan*. 1989 Jan 1; 58(1): 59-70.
- ²⁵⁹ Nihei-H; Kanemitsu-H; Tamura-A; Oka-H; Sano-K. Cerebral uric acid, xanthine, and hypoxanthine after ischemia: the effect of allopurinol. *Neurosurgery*. 1989 Oct; 25(4): 613-7.
- ²⁶⁰ Neglen P; Jabs CM; Eklof B. Plasma metabolic disturbances and reperfusion injury following partial limb ischaemia in man. *Eur J Vasc Surg*; 1989 Apr; 3(2); P 165-72.
- ²⁶¹ Leibovitz-B; Hu-ML; Tappel-AL. Dietary supplements of vitamin E, beta-carotene, coenzyme Q10 and selenium protect tissues against lipid peroxidation in rat tissue slices. *J-Nutr*. 1990 Jan; 120(1): 97-104.
- ²⁶² Maderazo-EG; Woronick-CL; Hickingbotham-N; Mercier-E; Jacobs-L; Bhagavan-H. Additional evidence of autoxidation as a possible mechanism of neutrophil locomotory dysfunction in blunt trauma. *Crit-Care-Med*. 1990 Feb; 18(2): 141-7.
- ²⁶³ Sangha O, Stucki G. Vitamin E in the treatment of rheumatic diseases. *Zeitschrift für Rheumatologie* 1998;57(4):207-214.
- ²⁶⁴ Sacheck JM, Blumberg JB. Role of vitamin E and oxidative stress in exercise. *Nutrition* 2001;(10):809-14.
- ²⁶⁵ Zafarullah M, Li WQ, Sylvester J, Ahmad M. Molecular mechanisms of N-acetylcysteine actions. *Cell Mol Life Sci* 2003;60(1):6-20.
- ²⁶⁶ Andriamanalijaona R, Kypriotou M, Bauge C, Renard E, Legendre F, Raoudi M, Boumediene K, Gatto H, Monginoux P, Pujol JP. Comparative effects of 2 antioxidants, selenomethionine and epigallocatechin-gallate, on catabolic and anabolic gene expression of articular chondrocytes. *J Rheumatol*. 2005;32(10):1958-67.
- ²⁶⁷ Akimoto AK, Miranda-Vilela AL, Alves PC, Pereira LC, Lordelo GS, Hiragi Cde O, da Silva IC, Grisolia CK, Klautau-Guimarães Mde N. Evaluation of gene polymorphisms in exercise-induced oxidative stress and damage. *Free Radic Res*. 2010 Mar;44(3):322-31.
- ²⁶⁸ Miranda-Vilela AL, Lordelo GS, Akimoto AK, Alves PC, Pereira LC, Klautau-Guimarães MD, Grisolia CK. Genetic polymorphisms influence runners' responses to the dietary ingestion of antioxidant supplementation based on pequi oil (*Caryocar brasiliense* Camb.): a before-after study. *Genes Nutr*. 2011 Nov;6(4):369-95.